



DEPARTMENT OF
EDUCATION

**NATIONAL
SENIOR CERTIFICATE**

GRADE/GRAAD 12

**MATHEMATICS PAPER 2/ WISKUNDE V2
SEPTEMBER 2020
MEMORANDUM**

MARKS/PUNTE: 150

This memorandum consists of 15 pages / Hierdie memo bestaan uit 15 bladsye.


QUESTION /VRAAG 1

1.1.1		✓ box /mond ✓ whiskers, on scale/ stert, en op skaal (2)
1.1.2	<p>School B/skool B: In B 50% of learners got above 60% whilst 50% of learners of A above 55% / 50% van B het meer as 60% ; by A het 50% meer as 55% median of B > median of A/ mediaan van B > mediaan van A 75% of B above 30% and at A 75% above 23,5% / 75% van B bo 30% ; By A 75% bo 23,5%</p> <p>OR/OF Skewed to the right/ skeef na Regs</p>	✓ School /skool B ✓✓ any two reasons enige twee redes ✓✓ OR/ OF ✓✓ (3)
1.2	$\frac{135 \times 24 + 225 \times 32 + 200 \times x}{560} = x$ $3240 + 7200 + 200x = 560x$ $360x = 10440$ $x = 29$	✓ numerator/ teller ✓ 560 ✓ 360x = 10440 ✓ answer/antwoord (4)

QUESTION/VRAAG 2

2.1	$a = 23,19$ $b = 0,768..$ $y = 23,19 + 0,77x$	✓ a ✓ b ✓ $y = 23,19 + 0,77x$ (3)
2.2	$y = 23,19 + 0,77(30)$ $= 46,29$ $\approx 46\%$	✓ substitution/vervanging ✓ answer/antwoord (2)


2.3.1	$\delta = 10,99$	✓✓answer <i>/antwoord</i> (2)
2.3.2	$57,4\% - 49,4\% = 8\%$ $\frac{x}{50} = \frac{8}{100}$ $x = 4 \text{ marks}$	✓8% ✓4 marks/ punte (2) [9]

QUESTION /VRAAG 3

3.1	$x + ay = a$ $0 + ay = a$ $y = 1$ $\therefore P(0;1)$	✓x=0 ✓y=1 ✓P(0;1) (3)
3.2	$OS : OP = 3 : 1$ $S(3;0) : x + ay = a$ $3 + a(0) = a$ $a = 3$	✓S(3;0) ✓substitution/vervang ✓a = 3 (3)
3.3	$m_{PS} = -\frac{1}{3}$ $m_{TR} = 3$ RT: $y = 3x - 5\frac{2}{3}$	✓ $m_{PS} = -\frac{1}{3}$ ✓ $m_{TR} = 3$ ✓equation/ vergelyking (3)
3.4	$x + 3y = 3 \Rightarrow y = -\frac{1}{3}x + 1$ $\therefore -\frac{1}{3}x + 1 = 3x - \frac{17}{3}$ $-x + 3 = 9x - 17$ $20 = 10x$	✓equating /stel gelyk

	<p>Substitute: $y = -\frac{1}{3}(2) + 1$ $y = \frac{1}{3}$ $R(2; \frac{1}{3})$</p>	<p>✓ simplification/vereenvoudig ✓ $x = 2$ ✓ $y = \frac{1}{3}$ (4)</p>
3.5	$\begin{aligned} \text{area} \Delta PRT &= \frac{1}{2} \times PT \times h \\ &= \frac{1}{2} \times 6 \frac{2}{3} \times 2 \\ &= \frac{20}{3} \text{ sqd units/ vierkante eenhede} \end{aligned}$	<p>✓ $PT = 6 \frac{2}{3}$ ✓ $h = 2$ ✓ answer/ antwoord (3)</p>
3.6	<p>PT is the diameter (converse: \angle in semi \odot)/ $\odot PT$ is die middellyn (omgekeerde: \angle in semi \odot) $\text{radius} = \frac{10}{3}$ units.</p>	<p>✓ S/R ✓ answer / antwoord (2)</p>
		[18]


QUESTION/VRAAG 4




4.1.1	$x^2 - 6x + 9 + y^2 - 4y + 4 = 12 + 9 + 4$ $(x - 3)^2 + (y - 2)^2 = 25$ $C(3;2)$ $r = 5$	Completing the square <i>vierkantsvoltooiing</i> ✓ LHS/ <i>LK</i> ✓ RHS/ <i>RK</i> ✓ $C(3;2)$ ✓ $r = 5$ (4)
4.1.2	Y-intercept: /Y as: $x = 0$ $y^2 - 4y - 12 = 0$ $(y - 6)(y + 2) = 0$ $y = -2 \text{ or } y = 6 \therefore A(0;6)$ $m_{AC} = \frac{6-2}{0-3} = -\frac{4}{3}$ $m_{AB} = \frac{3}{4}$ (tan \perp radius) $y = \frac{3}{4}x + 6$	✓ $x = 0$ ✓ standardform/ <i>standaardvorm</i> ✓ factors/ <i>faktore</i> ✓ $A(0;6)$ ✓ $m_{AC} = -\frac{4}{3}$ ✓ $m_{AB} = \frac{3}{4}$ ✓ R ✓ equation/ <i>vergelyking</i> (8)
4.1.3	Let $A = \theta$: $\tan \theta = \frac{3}{4}$ $\theta = 36,87^\circ$ $\alpha = 90^\circ + 36,87^\circ$ $\alpha = 126,87^\circ$	✓ $\tan \theta = \frac{3}{4}$ ✓ $\theta = 36,87^\circ$ ✓ $\alpha = 90^\circ + 36,87^\circ$ ✓ answer / <i>antwoord</i> (4)

4.2	$x^2 + 2x + 1 + y^2 - 4y + 4 = 44 + 1 + 4$ $(x+1)^2 + (y-2)^2 = 49$ Centre / middelpunt D(-1;2) ; radius = 7 $CD = 3 - 1 = 4$ $r_C + r_D = 5 + 7 = 12 > 4$ \therefore the two circles intercept internally/ <i>twEE sirkels sny inwendig</i>	✓Centre D(-1;2) ✓r = 7 ✓CD = 4 ✓ $r_C + r_D = 12$ ✓>4 ✓conclusion / <i>gevolgtrekking (6)</i> [22]


QUESTION/VRAAG 5

5.1	$\tan A = \frac{5}{12}$  $r^2 = 12^2 + 5^2$ $= 169$ $r = 13$ $13\sin A - 3\tan B$ $= 13\left(\frac{5}{13}\right) - 3\left(\frac{12}{5}\right)$ $= 5 - \frac{36}{5}$ $= -\frac{11}{5}$	✓diagram ✓r = 13 ✓ $\sin A = \frac{5}{13}$ ✓ $\tan B = \frac{12}{5}$ ✓answer /antwoord (5)
5.2	$\begin{aligned} & \frac{-\tan 60^\circ - \sin x}{\sin x + 2\cos 30^\circ} \\ &= \frac{-\sqrt{3} - \sin x}{\sin x + 2\left(\frac{\sqrt{3}}{2}\right)} \\ &= \frac{-(\sin x + \sqrt{3})}{(\sin x + \sqrt{3})} \\ &= -1 \end{aligned}$	✓ $-\tan 60^\circ$ ✓ $-\sin x$ ✓ $2\cos 30^\circ$ ✓special angles/ <i>bekende hoeke</i> ✓factorisation/ <i>faktoriseer</i> ✓answer/


5.3.1	$\begin{aligned} \cos(x+y) &= \sin(90^\circ - (x+y)) \\ &= \sin((90^\circ - x) + (-y)) \\ &= \sin(90^\circ - x)\cos(-y) + \cos(90^\circ - x)\sin(-y) \\ &= \cos x \cos y - \sin x \sin y \end{aligned}$	<ul style="list-style-type: none"> ✓ Correct co-fn/ korrekte ko-funksie ✓ regrouping/ herrangskik ✓ Correct expansion/ uitbreiding ✓ simplification/ vereenvoudig <p>(4)</p>
5.3.2	$\begin{aligned} \cos(x-y) - \cos(x+y) &= \cos x \cos y + \sin x \sin y - (\cos x \cos y - \sin x \sin y) \\ &= \cos x \cos y + \sin x \sin y - \cos x \cos y + \sin x \sin y \\ &= 2 \sin x \sin y \end{aligned}$	<ul style="list-style-type: none"> ✓ Expansion/ brei uit ✓ simplification / vereenvoudig <p>(2)</p>
5.3.3	$\begin{aligned} 2 \sin 195^\circ \sin 45^\circ &= \cos(195^\circ - 45^\circ) - \cos(195^\circ + 45^\circ) \\ &= \cos 150^\circ - \cos 240^\circ \\ &= -\cos 30^\circ - (-\cos 60^\circ) \\ &= -\frac{\sqrt{3}}{2} + \frac{1}{2} \\ &= \frac{-\sqrt{3} + 1}{2} \end{aligned}$	<ul style="list-style-type: none"> ✓ Substitution/ vervang ✓ $-\cos 30^\circ$ ✓ $-\cos 60^\circ$ ✓ Special angles/ bekende hoeke ✓ answer/antwoord <p>(5)</p>
		[22]

QUESTION/VRAAG 6

6.1	g : ✓ x-intercepts / x - asse ✓ y-intercepts / y - as ✓ turningpoints / draaipunte	(3)
6.2	$2\sin x = \cos x \cos 30^\circ - \sin x \sin 30^\circ$ $2\sin x = \frac{\sqrt{3}}{2} \cos x - \frac{1}{2} \sin x$ $\frac{5}{2} \sin x = \frac{\sqrt{3}}{2} \cos x$ $\frac{\sin x}{\cos x} = \frac{\sqrt{3}}{5}$ $\tan x = \frac{\sqrt{3}}{5}$	✓ Expansion of g / brei uit ✓ Special angles/ bekende hoekte ✓ $\frac{5}{2} \sin x = \frac{\sqrt{3}}{2} \cos x$ ✓ $\frac{\sin x}{\cos x}$
6.3	$\tan x = \frac{\sqrt{3}}{5}$ $x = 19,1^\circ \text{ or } x = 180^\circ + 19,1^\circ = 199,1^\circ$	✓ $19,1^\circ$ ✓ $199,1^\circ$ (2)
6.4.1	$x \in [60^\circ; 180^\circ] \cup [240^\circ; 270^\circ]$	✓ 60° and / en 180° ✓ 240° and / en 270° ✓ notation / notasie
6.4.2	$x \in [19,1^\circ; 199,1^\circ]$	✓ critical values / kritieke waardes ✓ notation / notasie (2)


QUESTION /VRAAG 7

$\text{In } \triangle ABC : \tan x = \frac{h}{CB}$ $CB = \frac{h}{\tan x}$ $\text{In } \triangle BCD : CB = BD$ $CD^2 = BC^2 + BD^2 - 2BC \times BD \times \cos y$ $=$ $(\frac{h}{\tan x})^2 + (\frac{h}{\tan x})^2 - 2 \frac{h}{\tan x} \times \frac{h}{\tan x} \times \cos y$ $= \frac{2h^2}{\tan^2 x} - \frac{2h^2 \cos y}{\tan^2 x}$ $= \frac{2h^2}{\tan^2 x} (1 - \cos y)$ $CD = \frac{h}{\tan x} \sqrt{2(1 - \cos y)}$ $CD = \frac{h}{\tan x} \sqrt{2 - 2 \cos y}$	✓ $\tan x = \frac{h}{CB}$ ✓ $CB = \frac{h}{\tan x}$ ✓ Correct application of cos rule/ korrekte gebruik van cosreël ✓ substitution ✓ simplify/ vereenvoudig ✓ factorising/ faktoriseer
	[6]

QUESTION/VRAAG 8

8.1.1	$\hat{Q}_2 = \hat{R}_2$ (tangents from same point) / raaklyne uit dsde punt) $2\hat{R}_2 + 70^\circ = 180^\circ$ ($\angle's$ of ΔPQR / $\angle'e$ van ΔPQR) $2\hat{R}_2 = 110^\circ$ $\hat{R}_2 = 55^\circ$	✓S✓R ✓S ✓answer/ antwoord (4)
8.1.2	$\hat{Q}_1 + \hat{Q}_2 = 90^\circ$ (tangent \perp radius) / (raaklyn \perp op radius) $\hat{Q}_1 + 55^\circ = 90^\circ$ $\hat{Q}_1 = 35^\circ$	✓S✓R ✓answer / antwoord (3)
8.1.3	$\hat{Q}_1 = \hat{R}_1 = 35^\circ$ (r =) $\hat{M}_1 = 110$ ($\angle's$ of ΔMQR) $\hat{M}_2 = \hat{M}_1 = 110$ (= chords subtend $=\angle's$ at centre)/ (= koorde = middelpunts $\angle'e$)	✓S/R ✓ $\hat{M}_2 = 110$ ✓R (3)


8.2



8.2.1	$\hat{X}_1 = 90^\circ$ (line from centre to midpoint of chord)/ $(van midpunt van \odot, mdpt van koord)$ $\hat{U}_3 + \hat{U}_4 = 90^\circ$ (\angle in semi \odot) $\therefore \hat{X}_1 = \hat{U}_3 + \hat{U}_4 = 90^\circ$ $\Rightarrow RU \parallel SV$ (corresponding \angle 's =)/ $(ooreenkomsige \angle'e =)$	$\checkmark S \checkmark R$ $\checkmark S$ $\checkmark S$ $\checkmark R$ (5)
8.2.2	$\hat{R}_2 = y$ (tan-chord theorem)/ $(\angle tussen raaklyn en koord)$ $\hat{O}_1 = y$ (alternate \angle 's, $RU \parallel SV$)/ $(verwisselende \angle'e, RU \parallel SV)$ $\hat{O}_1 = 2\hat{T}_1$ (\angle at centre = $2\angle$ at circumference) $(middelpunts\angle = 2 \times omtreks\angle)$ $\therefore \hat{T}_1 = \frac{1}{2}y$	$\checkmark S \checkmark R$ $\checkmark S$ $\checkmark S \checkmark R$ (5)
		[20]


QUESTION/VRAAG 9

9.1




9.1	<p>Construction: Draw altitudes h and k. Konstruksie: trek hoogtelyne h en k.</p> $\frac{\text{area}\triangle ADE}{\text{area}\triangle BDE} = \frac{\frac{1}{2} \times AD \times h}{\frac{1}{2} \times BD \times h} = \frac{AD}{DB}$ $\frac{\text{area}\triangle ADE}{\text{area}\triangle DEC} = \frac{\frac{1}{2} \times AE \times k}{\frac{1}{2} \times EC \times k} = \frac{AE}{EC}$ <p>But the $\text{area}\triangle BDE = \text{area}\triangle DEC$ (same base and same height/ <i>dsde basis en hoogte</i>)</p> $\therefore \frac{\text{area}\triangle ADE}{\text{area}\triangle BDE} = \frac{\text{area}\triangle ADE}{\text{area}\triangle DEC}$ $\therefore \frac{AD}{DB} = \frac{AE}{EC}$	<p>✓ Construction/ <i>konstruksie</i></p> <p>✓S</p> <p>✓S</p> <p>✓S✓R</p> <p>✓S</p>
		(6)

9.2



9.2.1	$\frac{AE}{EM} = \frac{AF}{FD} = \frac{4}{3}$ (line to one side of ΔAMD) $\frac{EM}{AM} = \frac{3}{7}$	$\checkmark S \checkmark R$ \checkmark answer (3)
9.2.2	$AM : MC = 7 : 7$ (diagonals of parm) $\frac{CM}{ME} = \frac{7}{3}$	$\checkmark S \checkmark R$ \checkmark answer (3)
9.2.3	$\text{area } \Delta BDC = \text{area } \Delta ABD$ (diagonals of parm) $\frac{\text{area } \Delta FDC}{\text{area } \Delta BDC} = \frac{\text{area } \Delta FDC}{\text{area } \Delta ABD}$ $= \frac{\frac{1}{2} \times FD \times \text{height}}{\frac{1}{2} \times AD \times \text{height}}$ $= \frac{FD}{AD}$ (between same parallel lines)/ $(tussen dieselfde \parallel lyne)$ $= \frac{3}{7}$	$\checkmark S$ \checkmark replacing/ <i>vervang</i> \checkmark area formula / <i>oppervlak formule</i> $\checkmark R$ (4)
		[16]

QUESTION / VRAAG 10



10.1	<p>In ΔXYC and ΔDYX:</p> $\hat{X}_1 = \hat{D}_1 \quad (\text{tan-chord theorem}/\angle \text{ tussen raaklyn en koord})$ $\hat{C}_2 = \hat{X}_2 \quad (\text{tan-chord theorem}/\angle \text{ tussen raaklyn en koord})$ $\hat{Y}_3 + \hat{Y}_4 = \hat{Y}_1 + \hat{Y}_2 (\angle's \text{ of } \Delta)$ $\therefore \Delta XYC \parallel \Delta DYX (\angle \angle \angle)$ $\Rightarrow \frac{XY}{DY} = \frac{YC}{YX}$ $\therefore XY^2 = YC \cdot DY$	$\checkmark S \checkmark R$
10.2	$\hat{A}_1 = 2\hat{X}_1 \quad (\angle \text{ at centre is twice } \angle \text{ at circumference})$ $(\text{middelpunts}\angle = 2 \times \text{omtreks}\angle)$ <p>Similarly/netso $\hat{B}_1 = 2\hat{D}_1$</p> <p>But $\hat{X}_1 = \hat{D}_1$ (tan-chord theorem/ $\angle \text{ tussen raaklyn en koord}$)</p> $\therefore \hat{A}_1 = \hat{B}_1$	$\checkmark S \checkmark R$
10.3	$\frac{CA}{YB} = \frac{CY}{YX} \quad (\Delta CAY \parallel \Delta YBX)$ $\frac{R}{r} = \frac{CY}{YX}$ $\therefore \frac{R^2}{r^2} = \frac{CY^2}{YX^2}$ $= \frac{CY \cdot CY}{CY \cdot DY}$ $= \frac{CY}{DY}$	$\checkmark S$ $\checkmark \text{Substitute/ vervang}$ $\checkmark \text{Square both sides/}$ $\text{kwadreer albei kante}$

	$\therefore \frac{r^2}{R^2} = \frac{DY}{CY}$	<p>✓Replace/vervang YX^2</p> <p>✓answer /antwoord</p>
		(5) [14]