

**GAUTENG PROVINCE**  
EDUCATION  
REPUBLIC OF SOUTH AFRICA

**PREPARATORY EXAMINATION  
2015  
MEMORANDUM**

**SUBJECT:**

**MATHEMATICS P1 (10611)**

GAUTENG DEPARTMENT OF EDUCATION  
PREPARATORY EXAMINATION – 2015

MATHEMATICS  
(First Paper)

**MEMORANDUM**

<b>QUESTION 1 [21 MARKS]</b>		
1.1	1.1.1	$x = -\frac{5}{2}$ or $x = \sqrt{2}$ or $x = -\sqrt{2}$
	1.1.2	$x^2 \geq 9$ or $x^2 - 9 \geq 0$ $x \geq 3$ or $x \leq -3$
		[3] 1A 1A for critical value of 3 2A for each inequality Max 2/3 if ‘and’ is used Full marks if ‘or’ is omitted
	1.1.3	$6^{2x} \cdot 2^{2x} = 8 \cdot 6^{2x}$ or $2^{4x} \cdot 3^{2x} = 8 \cdot 3^{2x} \cdot 2^{2x}$ or $2^{4x} \cdot 3^{2x} = 2^{3+2x} \cdot 3^{2x}$ $2^{2x} = 8$ $2x = 3$ $x = \frac{3}{2}$
		[4] 1A for factorizing bases 1CA 1CA 1CA
1.2		$(x+1)^2 = \frac{9}{2}$ $x+1 = \pm \sqrt{\frac{9}{2}}$ $x = 1, \frac{1}{2}$ or $x = -3, \frac{1}{2}$
		[4] 1A 1CA 2CA for each solution
		OR
		$2x^2 + 4x - 7 = 0$
		$x = \frac{-4 \pm \sqrt{4^2 - 4(2)(-7)}}{2(2)}$
		1A standard form 1CA substitution into correct formula 2CA for each solution
		[4]

1.3	$\frac{(-2x+7)+5}{x-1} = \frac{1}{2}$	1A correct substitution
	$2(-2x+12) = x-1$	1CA simplification
	$x = 5$	1CA
	Substitute $x = 5$ : $y = -2(5) + 7$	
	$y = -3$	1CA
	OR	
	$\frac{y+5}{\frac{y}{2} + \frac{7}{2} - 1} = \frac{1}{2}$	1A correct substitution
	$2(y+5) = -\frac{y}{2} + \frac{5}{2}$	1CA simplification
	$y = -3$	1 CA
	substitute $y = -3$ : $x = -\frac{-3}{2} + \frac{7}{2}$	
	$x = 5$	1CA
		[4]
1.4	$1-p > 0$	1A
	$p < 1$	1CA [2]

## QUESTION 2 [12 MARKS]

2.1	2.1.1	$x\left(-\frac{x}{2}\right)$	1A for $-\frac{x}{2}$
		$= -\frac{x^2}{2}$	1CA for multiplication with $x$
	2.1.2	$x+x+2$	[2]
		$= 2x+2$	1A for $x+2$
			1CA for adding to $x$
2.2		$-2+x+2x+2 = -\frac{x^2}{2}$	[2]
		$x^2+6x=0$	1M
		$x(x+6)=0$	1CA
		$x=-6$	1CA factorizing
			1CA choosing correct $x$ -value
2.3		The series does NOT converge.	[5]
		$r = 3$ . to converge, $-1 < r < 1$	2M for any logical explanation
			[3]

**QUESTION 3 [8 MARKS]**

3.1	$(250)^2 + 4(250)$ = 63500	1A substitution 1CA [2]
3.2	$T_{100} = S_{100} - S_{99}$ = $100^2 + 4(100) - [99^2 + 4(99)]$ = 203	1M 1A correct substitution 1CA
	OR	
	$T_1 = 5 ; T_2 = 7 ; T_3 = 9$	1A for first three terms
	$T_{100} = 5 + 99(2)$ = 203	1M for correct formula 1CA [3]
3.3	$n^2 + 4n = 1440$ $(n+40)(n-36) = 0$	1A 1CA correct factors or correct substitution into correct formula.
	$n = 36$	1CA choice of correct $n$ . Full marks for answer only [3]

**QUESTION 4 [6 MARKS]**

4.1	2; 6; 12; 20	1A for first three terms 1A for the 4 <sup>th</sup> term [2]
4.2	$a = 1; b = 1; c = 0$	3A finding $a, b, c$ using any correct method.
	$T_n = n^2 + n$ or $T_n = n(n+1)$	
	$T_{100} = 100(100+1)$ = $10100 \text{ cm}^2$	1CA [4]


**QUESTION 5 [15 MARKS]**

5.1	5.1.1	$A = 650\ 000(1 - 0,3)^4$ = R156 065	1A substitution into correct formula 1CA [2]
	5.1.2	$A = 650\ 000(1 + 0,15)^4$ = R1 136 854,06	1A substitution into correct formula 1CA [2]
	5.1.3	$1136854,06 - 156065 = 980789,06$ $980789,06 = \frac{x \left[ \left(1 + \frac{0,095}{12}\right)^{48} - 1 \right]}{0,095}$ = R16 875,92	1CA 1M for use of correct formula 1CA us of correct formula 1CA [4]
5.2	5.2.1	$OB = 1275\ 000 \left(1 + \frac{0,092}{12}\right)^{84} - \frac{11636,02 \left[ \left(1 + \frac{0,092}{12}\right)^{84} - 1 \right]}{0,092}$ = R1056 675,39	1M 1A correct substitution 1CA [3]
	5.2.2	$1056675,39 \left(1 + \frac{0,092}{12}\right)^5$ = R1097807,15	1CA substitution into correct formula 1CA 1CA substitution into correct formula 1CA [4]

**QUESTION 6 [3 MARKS]**

			1M for correct shape 1A for horizontal asymptote 1A for negative y-value [3]
--	--	--	---

QUESTION 7 [19 MARKS]		
7.1	$-2x^2 - 4x + 30 = 0$ $(x+5)(x-3) = 0$	1M equating to 0 1A correct factors
	$A(-5;0) \quad B(3;0)$	2CA for both intercepts -1 if not in coordinate form
		[4]
7.2	$y = -2(x^2 + 2x - 15)$ $y = -2(x^2 + 2x + 1 - 15 - 1)$	1A for taking out -2 1CA $\left(\frac{b}{2}\right)^2$ added and deducted
	$y = -2[(x+1)^2 - 16]$ $y = -2(x+1)^2 + 32$	1CA in complete square form 1CA
	Tpt $(-1;32)$	1CA
	OR	
	$x = \frac{-4}{2(-2)}$	1A substitute into correct formula
	$x = -1$	1CA
	$y = 32$	1CA
	$y = -2(x+1)^2 + 32$	1CA in correct form
	Tpt $(-1;32)$	1CA
		[5]
7.3	$y = -2(1)^2 - 4(1) + 30$	1M substitution into equation
	$y = 24$ - No	1CA value and conclusion
	OR	
	$-2x^2 - 4x + 30 = 2x + 10$	1M finding point of intersection
	$x^2 + 3x - 10 = 0$	
	$x = 2$ or $x = -5$ - No	1CA for values with conclusion
		[2]
7.4	$-2x^2 - 4x + 30 = mx + 32$	1M for equating
	$-2x^2 + (-4-m)x - 2 = 0$	1CA for standard form
	$\Delta = 0$	1M
	$(-4-m)^2 - 4(-2)(-2) = 0$	1CA correct substitution
	$m = 0$ or $m = -8$	2 CA for each value of $m$
		[6]
7.5	$(-\infty; 3) - \{-5\}$ or $x < 3; x \neq -5$	2A
		[2]

QUESTION 8 [11 MARKS]		
8.1	VA: $x = 2$	1A
	HA: $y = -1$	1A
		[2]
8.2	y-intercept: $y = \frac{1}{0-2} - 1$ $= -\frac{3}{2}$	1M 1CA
	$x$ -intercept: $\frac{1}{x-2} - 1 = 0$ $x = 3$	1M 1CA
		[4]
8.3		1A intercepts with axes 1A asymptotes 1A Shape [3]
8.4	$2 = \frac{1}{k-2} - 1$	1A correct substitution
	$k = \frac{4}{3}$	1CA
		[2]

QUESTION 9 [6 MARKS]		
9.1	$-2 = \log_a \left( \frac{9}{4} \right)$	1A correct substitution
	$a^{-2} = \left( \frac{9}{4} \right)$	1CA exponential form
	$a = \frac{2}{3}$	1CA
		[3]
9.2	$y = -\log_2 \frac{x}{3}$ or $y = \log_3 \frac{x}{2}$	1A [1]
9.3	$g(x) > -2$	1A [1]
9.4	$(-\infty; \infty)$	1A [1]


QUESTION 10 [14 MARKS]		
10.1	$\frac{f(3+h) - f(3)}{h}$	1M correct formula
	$= \frac{-2(3+h)^2 + 1 - [-2(3)^2 + 1]}{h}$	1A correct substitution
	$= \frac{-12h - 2h^2}{h}$	1CA correct simplification
	$= \frac{h(-12 - 2h)}{h}$	1CA
	$= -12 - 2h$	Max 1/3 if 'limit' is used
	OR	
	$\frac{f(x+h) - f(x)}{h}$	1M correct formula
	$= \frac{-2(x+h)^2 + 1 - [-2x^2 + 1]}{h}$	1A correct substitution
	$= \frac{-4xh - 2h^2}{h}$	1CA correct simplification
	$= \frac{h(-4(3) - 2h)}{h}$	1CA substitution of $x=3$
	$= -12 - 2h$	Max 1/3 if 'limit' is used
		[4]
10.1.2	$\lim_{h \rightarrow 0} (12 - 2h)$	1M using limit
	$= 12$	1CA
		[2]
10.1.3	$m = 0$	1A
		[1]

10.2	10.2.1	$y = 16 - 16x + 4x^2$	1A for simplification
		$y' \text{ or } \frac{dy}{dx} = -16 + 8x$	2CA (-1 for incorrect notation)
			[3]
	10.2.2	$y = x - ax^{\frac{1}{2}} + a$	1A for simplification
		$y' = 1 - \frac{1}{2}ax^{-\frac{1}{2}}$	2CA
		$y' = 1 - \frac{a}{2x^{\frac{1}{2}}}$	1CA positive exponents
			[4]

**QUESTION 11 [11 MARKS]**

11.1		The graph passes through the origin. Or $f(x) = (x - 0)^2(x - a)^2$	1M any logical explanation
			[1]
11.2	11.2.1	$f(x) = x(x - 2)^2$	1A factorization
		$a = 2$	1CA
			[2]
	11.2.2	$f'(x) = 0$	1M
		$3x^2 - 8x + 4 = 0$	1A correct differentiation
		$(3x - 2)(x - 2) = 0$	1CA factors
		$b = \frac{2}{3}$	1CA
			[4]
11.3		$f'(0) = 3(0)^2 - 8(0) + 4$	1CA substitution into derivative
		$m = 4$	1CA
			[2]
11.4		$x^3 - 4x^2 + 4x = p - 2$	1M taking 2 to RHS
		$p < 2$	1CA (full marks for answer only)
			[2]

QUESTION 12 [9 MARKS]		
12.1	$x^2 \left( 40 - \frac{x}{3} \right) < 0$	1M for inequality
	$x > 120 \text{ km/hour}$	1A for inequality sign 1CA for $120 \text{ km/hour}$ Full marks if units are omitted.
12.2		[3]
	$P = 40x^2 - \frac{x^3}{3}$	
	$P'(x) = 0$	1M
	$80x - x^2 = 0$	1CA
	$x(80 - x) = 0$	1CA correct factorization
	$x = 80 \text{ km/hour}$	1CA
	$P = \frac{40}{12} (80)^2 - \frac{80^3}{12 \cdot 3}$	1CA
	<del><math>P = 21333,33 \text{ Rand/day}</math></del> <del><math>\cancel{R} 85333,33</math></del>	1CA Full marks if units are omitted.
		[6]

QUESTION 13 [15 MARKS]			
13.1		$\frac{1}{6} + \frac{7}{16}$ $= \frac{29}{48}$	2A for both outcome 1M for addition 1CA for simplification
			[4]
13.2	13.2.1		3A for each section
	13.2.2	$x + \frac{1}{6} + \frac{5}{12} + \frac{1}{3} = 1$ $x = \frac{1}{12}$ $P(A) = \frac{1}{6} + \frac{1}{12} = \frac{1}{4}$	[3] 1M 1A 1CA
13.3		36 000	1A
			[1]
13.4	13.4.1	2.9!	2A
			[2]
	13.4.2	8.8!	2A
			[2]

