



Basic Education

**KwaZulu-Natal Department of Education
REPUBLIC OF SOUTH AFRICA**

MATHEMATICS P1

PREPARATORY EXAMINATION

SEPTEMBER 2016

MEMORANDUM


**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

MARKS: 150

This memorandum consists of 12 pages.

QUESTION 1

1.1.1	$2x(3x-5) = 0$ $x = 0 \quad \text{or} \quad x = \frac{5}{3}$	AA✓✓ Answers	(2)
1.1.2	$x^2 - 3x = 7$ $x^2 - 3x - 7 = 0$ $x = \frac{-(-3) \pm \sqrt{(-3)^2 - 4(1)(-7)}}{2(1)}$ $x = 4,54 \quad \text{or} \quad x = -1,54$	A✓ equation in std. form CA✓ substitution CACACACA✓✓ answers	(4)
1.1.3	$2x - 5\sqrt{x} = 3$ $2x - 3 = 5\sqrt{x}$ $4x^2 - 12x + 9 = 25x$ $4x^2 - 37x + 9 = 0$ $(4x-1)(x-9) = 0$ $x = \frac{1}{4} \quad \text{or} \quad x = 9$ <i>n/a</i>	A✓ isolating $5\sqrt{x}$ M✓ squaring both sides CA✓ standard form CA✓ factors CA✓ both answers CA✓ rejecting	(6)
	OR $2x - 5x^{\frac{1}{2}} - 3 = 0$ $\left(2x^{\frac{1}{2}} + 1\right)\left(x^{\frac{1}{2}} - 3\right) = 0$ $x^{\frac{1}{2}} = \frac{-1}{2} \quad \text{or} \quad x^{\frac{1}{2}} = 3$ No solution $x = 9$	A✓ standard form CA✓ factors CA✓ $x^{\frac{1}{2}} = \frac{-1}{2}$ CA✓ $x^{\frac{1}{2}} = 3$ CA✓ answer CA✓ rejecting	(6)
1.1.4	$2^x(3x+1) < 0$ $2^x > 0 \quad \text{for all } x \in R$ $\therefore 3x+1 < 0$ $x < -\frac{1}{3}$	A✓ $2^x > 0 \quad \text{for all } x \in R$ A✓ $3x+1 < 0$ A✓ $x < -\frac{1}{3}$	(3)
	OR [Graphically] 	A✓ sketch of $y = 2^x$ A✓ sketch of $y = 3x+1$ A✓ $x < -\frac{1}{3}$	(3)

1.2	$\begin{aligned} & 2^{99} \cdot 2^1 - 2^{99} \\ & 2^{99}(2 - 1) \\ & = 2^{99} \end{aligned}$	M✓ for taking 2^{99} as a common factor A✓ (2 - 1) A✓ answer	(3)
1.3	$\begin{aligned} & 2x - y = 3 \rightarrow (1) \\ & x^2 + 5xy + y^2 = 15 \rightarrow (2) \\ & \text{From (1): } y = 2x - 3 \\ & \text{Subst. (1) into (2)} \\ & x^2 + 5x(2x - 3) + (2x - 3)^2 = 15 \\ & x^2 + 10x^2 - 15x + 4x^2 - 12x + 9 = 15 \\ & 15x^2 - 27x - 6 = 0 \\ & (5x + 1)(3x - 6) = 0 \\ & x = \frac{-1}{5} \quad \text{or} \quad x = 2 \\ & y = \frac{-17}{5} \quad \text{or} \quad y = 1 \end{aligned}$ <p>OR</p> $\begin{aligned} & 2x - y = 3 \rightarrow (1) \\ & x^2 + 5xy + y^2 = 15 \rightarrow (2) \\ & \text{From (1): } x = \frac{y+3}{2} \\ & \text{Subst. (1) into (2)} \\ & \left(\frac{y+3}{2}\right)^2 + 5y\left(\frac{y+3}{2}\right) + y^2 = 15 \\ & \left(\frac{y^2 + 6y + 9}{4}\right) + \frac{5y^2 + 15y}{2} + y^2 = 15 \\ & y^2 + 6y + 9 + 10y^2 + 30y + 4y^2 = 60 \\ & 15y^2 + 36y - 51 = 0 \\ & 5y^2 + 12y - 17 = 0 \\ & (5y + 17)(y - 1) = 0 \\ & y = \frac{-17}{5} \quad \text{or} \quad y = 1 \\ & x = \frac{-1}{5} \quad \text{or} \quad x = 2 \end{aligned}$	A✓ y as subject CA✓ substitution $y = 2x - 3$ CA✓ std. form CA✓ factors CA✓ x values CA✓ y values	(6)
			[24]

QUESTION 2

2.1	$2a = 4$ $a = 2$ $3a + b = 0$ $b = -6$ $a + b + c = 4$ $c = 8$ $T_n = 2n^2 - 6n + 8$	A✓ a value CA✓ b value CA✓ c value CA✓ general term	(4)
	OR		
	$2a = 4$ $a = 2$ $3a + b = 0$ $b = -6$ $T_0 = c = 8$ $T_n = 2n^2 - 6n + 8$	A✓ a value CA✓ b value CA✓ c value CA✓ general term	(4)
	OR		
	$\begin{aligned} T_n &= T_1 + (n-1)d_1 + \frac{(n-1)(n-2)}{2}d_2 \\ &= 4 + (n-1)(0) + \frac{(n-1)(n-2)}{2}(4) \\ &= 5 + 0 + 2n^2 - 6n + 4 \\ &= 2n^2 - 6n + 8 \end{aligned}$	A✓ formula A✓ substituting first and second difference values CA✓ simplifying CA✓ general term	(4)
	OR		
2.2	$T_n = 4n - 4 = 28088$ $4n = 28092$ $n = 7023$ Between the 7023 rd and 7024 th terms.	A✓ $4n - 4$ CA✓ $4n - 4 = 28088$ CA(n ∈ N)✓ $n = 7023$ CA✓(n ∈ N)conclusion	(4)
	$T_n - T_{n-1} = 28088$ $2n^2 - 6n + 8 - [2(n-1)^2 - 6(n-1) + 8] = 28088$ $2n^2 - 6n + 8 - 2n^2 + 4n - 2 + 6n - 6 - 8 - 28088 = 0$ $4n - 28096 = 0$ $4n = 28096$ $n = 7024$ Between the 7023 rd and 7024 th terms.	A✓ $T_n - T_{n-1} = 28088$ CA✓ $4n - 28096 = 0$ CA✓(n ∈ N) $n = 7024$ CA✓(n ∈ N)conclusion	(4)
			[8]

QUESTION 3

3.1.1	$6 ; 2 ; 10 ; 2 ; 14 ; 2 ; \dots$ $18 ; 2$	AA✓✓ answers	(2)
3.1.2	$a = 2$ and is a constant sequence $\therefore S_{50} = 50 \times 2$ $= 100$	A✓100	(1)
3.1.3	$a = 6; d = 4; n = 50$ $S_{50} = \frac{50}{2} [2(6) + 4(50 - 1)]$ $= 5200$ $S_{100} = 5200 + 100$ $= 5300$	A✓ for a, d and n values CA✓ for substitution into formula CA✓ 5200 CA✓ answer	(4)
3.2	$S_n = a + ar + ar^2 + \dots + ar^{n-1} \rightarrow (1)$ $rS_n = ar + ar^2 + \dots + ar^n \rightarrow (2)$ (2) – (1): $rS_n - S_n = ar^n - a$ $S_n(r - 1) = a(r^n - 1)$ $\therefore S_n = \frac{a(r^n - 1)}{r - 1}$	A✓ for equation (1) A✓ for equation (2) A✓ subtraction on LHS and RHS A✓ factorising if $S_n = \frac{a(1 - r^n)}{1 - r}$ penalize 1 mark	(4)
			[11]

QUESTION 4

4.1	$a = 1 \quad r = \frac{2}{3}$ $S_\infty = \frac{a}{1-r}$ $= \frac{1}{1-\frac{2}{3}}$ $= 3$	A✓ a and r value CA✓ substitution of r into formula CA✓ answer	(3)
4.2	$S_n = 3n^2 - n$ $T_4 = S_4 - S_3$ $= [3(4)^2 - (4)] - [3(3)^2 - (3)]$ $= 44 - 24$ $= 20$ OR	M✓ $T_4 = S_4 - S_3$ AA✓✓ substitution A✓ answer	(4)

	$S_1 = T_1 = a = 2$ $S_2 = T_1 + T_2 = 3(2)^2 - 2 = 10$ $T_2 = 10 - 2 = 8$ $S_3 = T_1 + T_2 + T_3 = 3(3)^2 - 3 = 24$ $T_3 = 24 - 8 - 2 = 14$ $d = 6$ $T_4 = a + 3d = 2 + 3(6) = 20$	A✓ a value A✓ second term value CA✓ d value CA✓ answer	(4)
			[7]

QUESTION 5

5.1	$A = P(1-i)^n$ $A = 255000(1-0,125)^7$ $= R 100\ 137,46$	A✓ formula A✓ correct substitution CA✓ answer	(3)
5.2.1	$A = P(1+i)^n$ $= 10\ 000 \left(1 + \frac{0,095}{12}\right)^5$ $= R10\ 402,15$	A✓ correct substitution into correct formula CA✓ answer	(2)
5.2.2	$P = \frac{x[1 - (1+i)^{-n}]}{i}$ $10\ 402,15 = \frac{450 \left[1 - \left(1 + \frac{0,095}{12}\right)^{-n}\right]}{\frac{0,095}{12}}$ $\frac{10\ 402,15 \times \frac{0,095}{12}}{450} = 1 - \left(1 + \frac{0,095}{12}\right)^{-n}$ $\left(1 + \frac{0,095}{12}\right)^{-n} = 0,816999213$ $-n = \frac{\log 0,816999213}{\log \left(1 + \frac{0,095}{12}\right)}$ $n = 25,6315128$ $\therefore n = 26 \text{ payments}$	CA✓ correct substitution of P into formula A✓ $x = 450$ CA✓ n as subject M✓ using logs CA($n \in \mathbb{N}$)✓ answer	(5)

5.2.3	<p>Balance on Loan after the 25th payment</p> $= 10\ 402,15 \left(1 + \frac{0,095}{12}\right)^{25} - \frac{450 \left[\left(1 + \frac{0,095}{12}\right)^{25} - 1 \right]}{\frac{0,095}{12}}$ $= R12668,90 - R12386,54$ $= R282,36$ <p>OR</p> $P = \frac{x[1 - (1+i)^{-n}]}{i}$ $= \frac{450 \left[1 - \left(1 + \frac{0,095}{12}\right)^{-0,6315128} \right]}{\frac{0,095}{12}}$ $= R282,36$	<p>M✓ $(A - F_v)$ formula A✓ $n = 25$ CA✓ $10\ 402,15 \left(1 + \frac{0,095}{12}\right)^{25}$ CA✓ $\frac{450 \left[\left(1 + \frac{0,095}{12}\right)^{25} - 1 \right]}{\frac{0,095}{12}}$ CA✓ answer</p> <p>OR</p> <p>A✓ formula A✓ subst. 450 CA✓ subst $\frac{0,095}{12}$ CA✓ subst $0,6315128$ CA✓ answer</p>	(5)
			[15]

QUESTION 6 (For learners who did not receive the errata and used P(-4;4) , should be marked accordingly)

6.1	$p = -3$ $q = -2$	A✓ p value A✓ q value	(2)
6.2	$h(x) = \frac{a}{x+p} + q$ $P(4;-4) \text{ is a point on } h$ $-4 = \frac{a}{4-3} - 2$ $\therefore a = -2$	CA✓ subst. p, q and point P CA(negative)✓ a value	(2)
6.3	$h(x) = \frac{-2}{x-3} - 2$ $h(0) = \frac{-2}{0-3} - 2$ $= -1\frac{1}{3}$ $\left(0; -1\frac{1}{3}\right)$	A✓ substituting $x = 0$ CA(negative)✓ answer	(2)
6.4	$x = 1$	CACA✓✓ answer	(2)

6.5	$y = -(x + p) + q$ $y = -(x - 3) - 2$ $y = -x + 1$ $\therefore c = 1$ OR $y = -x + c$ <i>Point of intersection of asymptotes (3 ; -2)</i> $-2 = -3 + c$ $c = 1$	CA✓ substitution of p and q values into equation of line of symmetry CA✓ answer CA✓ substitution of $(3 ; -2)$ into equation of line of symmetry CA✓ answer	(2) (2)
			[10]

QUESTION 7

7.1	D(0 ; -10)	A(must be in coordinate form)✓ answer	(1)
7.2	$x^2 - 3x - 10 = 0$ $(x + 2)(x - 5) = 0$ $x = -2 \quad or \quad x = 5$ A(-2;0) B(5;0)	A✓ $x^2 - 3x - 10 = 0$ CA✓ factors CACAC(negative and positive)✓✓ each x -value A(-2;0) B(5;0)	(4)
7.3	$a = 2$ and $q = -10$	CA(positive)✓ a -value A✓ q -value	(2)
7.4	$x = -\frac{b}{2a} = -\frac{(-3)}{2(1)} = \frac{3}{2}$ or $x = \frac{-2+5}{2} = \frac{3}{2}$ $y = \left(\frac{3}{2}\right)^2 - 3\left(\frac{3}{2}\right) - 10 = -\frac{49}{4} / -12,25 / -12\frac{1}{4}$ C(1,5 ; -12,25)	A✓ $x = -\frac{b}{2a} = -\frac{(-3)}{2(1)} = \frac{3}{2}$ or CA✓ $x = \frac{-2+5}{2} = \frac{3}{2}$ CA✓ substitution CA✓ minimum value	(3)
7.5	(-1,5 ; -9,25)	CA✓ x -value CA✓ y -value	(2)
7.6	$x \geq \frac{3}{2}$ OR $g'(x).h'(x) \geq 0$ $(2x - 3).2 \geq 0$ $x \geq \frac{3}{2}$	CACAC✓✓ answer CA✓ product of derivatives CA(positive)✓ answer penalize 1 mark for incorrect notation	(2) (2)
			[14]

QUESTION 8

8.1	$y = \log_3 x$	AA✓✓ answer	(2)
8.2		A✓ Shape of p and p^{-1} A✓ y -intercept of p A✓ x -intercept of p^{-1} A✓ point on each graph	
8.3	$\log_3 x = 3$ $x = 27$ $0 < x \leq 27$	M✓ setting up equation CA✓ $x = 27$ CACA✓✓ for end points and inequality ANSWER ONLY full marks	(4)
			[10]

QUESTION 9 [Penalise once for notational error in questions 9, 10 and 11]

9.1	$f(x) = x^3$ $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ $= \lim_{h \rightarrow 0} \frac{(x+h)^3 - x^3}{h}$ $= \lim_{h \rightarrow 0} \frac{x^3 + 3x^2h + 3xh^2 + h^3 - x^3}{h}$ $= \lim_{h \rightarrow 0} \frac{h(3x^2 + 3xh + h^2)}{h}$ $= \lim_{h \rightarrow 0} (3x^2 + 3xh + h^2)$ $= 3x^2$	A✓ formula A✓ substitution into correct formula CA✓ simplifying CA✓ factors CA✓ answer	(5)
9.2.1	$f(x) = x^{\frac{1}{2}} - 4x^{-2}$ $f'(x) = \frac{1}{2}x^{-\frac{1}{2}} + 8x^{-3}$	AA✓✓ exponential form CACA✓✓ answer	(4)

<p>9.2.2</p> $\left(\frac{1}{2}x^2 - 3\right)^2 = y$ $y = \frac{1}{4}x^4 - 3x^2 + 9$ $\frac{dy}{dx} = x^3 - 6x$ <p>OR</p> $\frac{dy}{dx} = 2\left(\frac{1}{2}x^2 - 3\right) \cdot x$ $= x^3 - 6x$	<p>A✓ squaring both sides</p> <p>A✓ simplifying</p> <p>CACA✓✓ answers</p> <p>OR</p> <p>AA✓✓ derivative using chain rule</p> <p>CACA✓✓ answers</p>	<p>(4)</p> <p>(4)</p>
		<p>[13]</p>

QUESTION 10

10.1.1	$f(-3) = (-3)^2 - 8 = 1$	A✓ $y = 1$ (y - value)	(1)
10.1.2	$f'(x) = 2x$ $m = f'(-3) = 2(-3) = -6$	A✓ value of gradient of tangent	(1)
10.1.3	Equation of tangent :- $y = mx + c$ $1 = -6(-3) + c$ $c = -17$ $y = -6x - 17$ <p>OR</p> $y - 1 = -6(x + 3)$ $y = -6x - 18 + 1$ $y = -6x - 17$	CA✓ substitution CA✓ answer	(2)
10.2.1	$f(x) = x^3 - 3x - 2$ $f'(x) = 3x^2 - 3 = 0$ $3(x+1)(x-1) = 0$ $x = -1 \quad or \quad x = 1$ $y = 0 \quad or \quad y = -4$ $A(-1 ; 0) \quad B(1 ; -4)$	A✓ derivative <u>and</u> equating to 0 CA✓ factors CA✓ x - values CA✓ y - values	(4)

10.2.2	$ \begin{aligned} PQ &= g(x) - f(x) \\ &= (2x - 2) - (x^3 - 3x - 2) \\ &= -x^3 + 5x \\ PQ' &= -3x^2 + 5 = 0 \\ x^2 &= \frac{5}{3} \\ x &= \pm\sqrt{\frac{5}{3}} = 1,29 \\ \text{For } x &= 1,29 \\ y &= -(1,29)^3 + 5(1,29) = 4,3 \\ \text{Maximum Length of } PQ &= 4,3 \text{ units} \end{aligned} $	A✓ expression for PQ A✓ $PQ' = 0$ CA✓ x -values CA✓ y value for $x = 1,29$	(4)
10.2.3	$k = -4 \quad \text{or} \quad k = 0$	$k = 0$ A✓ $k = -4$ CA✓ answers	(2)
10.2.4	$f''(x) = 6x > 0$ $x > 0$ OR $x = \frac{x_1 + x_2}{2} = \frac{(-1) + (1)}{2} = 0$ $x > 0$ OR $x = -\frac{b}{3a} = -\frac{0}{3(1)} = 0$ $x > 0$	A✓ $6x$ A✓ $6x > 0$ A✓ answer A✓ midpoint formula CA✓ $x = 0$ CA✓ answer A✓ formula A✓ $x = 0$ A✓ answer	(3) (3) (3)
			[17]

QUESTION 11 (Ignore units in the question)

11.1	$V = s'(t) = 2t + 15 \text{ m/s}$	A✓ answer	(1)
11.2	$V = s'(25) = 2(25) + 15 = 65 \text{ m/s}$	CA✓ answer	(1)
11.3	$V = s'(t) = 2t + 15 \text{ m/s}$ $A = s''(t) = 2 \text{ m/s}^2$	CACA✓✓✓ 2	(2)
11.4	$A = s''(5) = 2 \text{ m/s}^2$	CA✓ answer (answer only 1 mark)	(1)
11.5	$s(t) = t^2 + 15t = 250$ $t^2 + 15t - 250 = 0$ $(t + 25)(t - 10) = 0$ $t = 10 \text{ s}$ $V = s'(10) = 2(10) + 15 = 35 \text{ m/s}$	A✓ equating s to 250 CA✓ factors CA✓ $t = 10$ CA✓ answer	(4)
			[9]

QUESTION 12

12.1.1	150	A✓ answer	(1)
12.1.2(a)	$P(\text{Male}) = \frac{70}{150} = \frac{7}{15}$	A✓ answer	(1)
12.1.2(b)	$P(\text{Eating Chocolate}) = \frac{80}{150} = \frac{8}{15}$	A✓ answer	(1)
12.1.3	$P(\text{Male eating chocolate}) = \frac{45}{150} = 0,3$ $P(\text{Male}) \times P(\text{Eating chocolate})$ $= \frac{70}{150} \times \frac{80}{150} = \frac{56}{225} = 0,249$ P(Male and Eating chocolate) ≠ P(Male) × P(Eating chocolate) Events are not independent	A✓ P(male eating chocolate) CA✓ P(Male) x P(eating chocolate) value CA✓ conclusion	(3)
12.2.1	$6 \times 7 \times 7 \times 7 = 2058$	A✓ $6 \times 7 \times 7 \times 7$ A✓ 2058	(2)
12.2.2	$6 \times 6 \times 5 \times 4 = 720$	A✓ $6 \times 6 \times 5 \times 4$ A✓ 720	(2)
12.2.3	Four digit codes divisible by 5:- $6 \times 7 \times 7 \times 1 = 252$ Probability of a four digit code divisible by 5 $= \frac{252}{2058} = \frac{7}{49} = 0,1428 = 14,28 \%$	A✓ $6 \times 7 \times 7 \times 1$ A✓ 252 CA✓ answer	(3)
			[13]

TOTAL MARKS: 150