



education

Department:
Education
PROVINCE OF KWAZULU-NATAL

**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

PHYSICAL SCIENCES: CHEMISTRY (P2)

COMMON TEST

MARCH 2019

MARKS : 50

TIME : 1 HOUR

This question paper consists of 6 pages and 2 data sheets.

INSTRUCTIONS AND INFORMATION TO CANDIDATES

1. Write your name and other information in the appropriate spaces on the ANSWER BOOK.
2. The question paper consists of FOUR questions. Answer ALL the questions in the ANSWER BOOK.
3. Start EACH question on a NEW page in the ANSWER BOOK.
4. Number the answers correctly according to the numbering system used in this question paper.
5. Leave one line between two sub-questions, for example between QUESTION 2.1 and QUESTION 2.2.
6. You may use a non-programmable pocket calculator.
7. You may use appropriate mathematical instruments.
8. You are advised to use the attached DATA SHEETS.
9. Show ALL formulae and substitutions in ALL calculations.
10. Round off your final numerical answers to a minimum of TWO decimal places where applicable.
11. Give brief motivations, discussions, et cetera where required.
12. Write neatly and legibly.

QUESTION 1: MULTIPLE – CHOICE QUESTIONS

Various options are provided as possible answers to the following questions. Choose the answer and write only the letter (A - D) next to the question number (1.1. – 1.4) in the ANSWER BOOK, for example 1.5 D.


1.1 Which ONE of the following compounds represented below is an aldehyde?

- A CH_3COOH
 B CH_3COCH_3
 C $\text{CH}_3\text{CH}_2\text{CHO}$
 D $\text{CH}_3\text{CH}(\text{OH})\text{CH}_3$ (2)

1.2 Boiling point is the . . .

- A temperature at which a liquid becomes a gas and is also called vapourisation point.
 B temperature at which a gas becomes a liquid and is also called condensation point
 C temperature at which a solid becomes a gas and is also called sublimation point
 D temperature at which the vapour pressure of a substance equals the atmospheric pressure. (2)


1.3 The structural formula of an organic compound is given below:



The correct IUPAC name for the above compound is . . .

- A 2,4 – dimethylhex – 1 – ene.
 B 3,5 – dimethylhex – 5 – ene.
 C 4 – ethyl – 2 – methylpent – 1 – ene.
 D 2 – ethyl – 4 – methylpent – 4 – ene. (2)

1.4 What are the reactants M and N in the following acid-catalysed reaction?




- A Methanoic acid and ethanol.
 B Ethanoic acid and methanol.
 C Methanoic acid and methanol.
 D Ethanoic acid and propan – 2 – ol. (2)

[8]

QUESTION 2 (Start on a new page.)

2.1 Consider the following compound:




2.1.1 Name the functional group of the above compound. (1)

2.1.2 Write down the IUPAC name of the above compound. (2)

2.1.3 Write down a structural formula of an isomer of the above compound. (2)

2.2 Consider the following organic compound:




2.2.1 Write down the general formula of the homologous series to which this compound belongs. (2)

2.2.2 Using MOLECULAR FORMULAE, write down a balanced equation for the reaction of the above compound with excess oxygen. (3)

2.3 A chemical analysis of a compound, $\text{C}_n\text{H}_{2n}\text{O}_2$, shows that it is made up of 12,5 % oxygen by mass. Calculate the percentage carbon in the compound. (5)

2.4 Polymers are organic compounds with a wide range of applications. The following compound represents a polymer formed as a result of addition polymerization:



2.4.1 Write down the structural formula of a monomer for the above. (2)

2.4.2 Write down the IUPAC name of the above polymer. (2)

[19]

QUESTION 3 (Start on a new page.)

An experiment was conducted to determine the boiling points of three(3) organic compounds, P, Q and R, from DIFFERENT HOMOLOGOUS SERIES under the same conditions. The following data was obtained:

COMPOUND	BOILING POINT/°C
P	103
Q	137
R	163


- 3.1 Define the term homologous series. (2)
- 3.2 Formulate an investigative question for this experiment. (2)
- 3.3 Which ONE of the compounds P, Q or R has the highest vapour pressure? Refer to the data in the table to give a reason for the answer. (2)
- 3.4 The three organic compounds used in the experiment are $\text{CH}_3(\text{CH}_2)_3\text{CH}_2\text{OH}$, $\text{CH}_3(\text{CH}_2)_2\text{COOH}$ and $\text{CH}_3(\text{CH}_2)_3\text{CHO}$, in no specific order.
- 3.4.1 Which ONE of these compounds is represented by letter R? (1)
- 3.4.2 Explain the answer to question 3.4.1 by referring to intermolecular forces and energy. (4)
- 3.5 How will the boiling point of pentane – 1,2 – diol compare to the boiling point of $\text{CH}_3(\text{CH}_2)_3\text{CH}_2\text{OH}$? (Choose from GREATER THAN, LESS THAN or EQUAL TO) (1)

[12]

Downloaded from Stanmorephysics.com

QUESTION 4 (Start on a new page.)

Three reactions of organic compounds are shown below:



4.1 Compound X represents an alcohol.

4.1.1 Name the type of reaction represented by reaction I. (1)

4.1.2 List any TWO reaction conditions that must be satisfied for reaction I to take place. (2)

4.1.3 Write down the IUPAC name for compound X. (2)

4.2 Reaction II is an elimination reaction.

4.2.1 Name the type of elimination reaction represented by reaction II. (1)

4.2.2 Name the inorganic product Y. (1)

4.2.3 Is the organic product of this reaction a MAJOR or MINOR product? (1)

4.3 In reaction III, an ALKENE Z is mixed with HCl.

4.3.1 Name this type of reaction. (1)

4.3.2 Explain why water must NOT be present in the reaction vessel. (2)

[12]

TOTAL MARKS: 50

**DATA FOR PHYSICAL SCIENCES GRADE 12
PAPER 2 (CHEMISTRY)**

**GEGEWENS VIR FISIESTE WETENSAPPE GRAAD 12
VRAESTEL 2 (CHEMIE)**

TABLE 1: PHYSICAL CONSTANTS/TABEL 1: FISIESTE KONSTANTES

NAME/NAAM	SYMBOL/SIMBOOL	VALUE/WAARDE
Standard pressure <i>Standaarddruk</i>	p°	$1,013 \times 10^5 \text{ Pa}$
Molar gas volume at STP <i>Molêre gasvolume by STD</i>	V_m	$22,4 \text{ dm}^3 \cdot \text{mol}^{-1}$
Standard temperature <i>Standaardtemperatuur</i>	T°	273 K
Charge on electron <i>Lading op elektron</i>	e	$-1,6 \times 10^{-19} \text{ C}$
Avogadro's constant <i>Avogadro-konstante</i>	N_A	$6,02 \times 10^{23} \text{ mol}^{-1}$

TABLE 2: FORMULAE/TABEL 2: FORMULES

$n = \frac{m}{M}$	$n = \frac{N}{N_A}$
$c = \frac{n}{V}$ or/of $c = \frac{m}{MV}$	$n = \frac{V}{V_m}$
$\frac{c_a V_a}{c_b V_b} = \frac{n_a}{n_b}$	$\text{pH} = -\log[\text{H}_3\text{O}^+]$
$K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1 \times 10^{-14}$ at/by 298 K	
$E_{\text{cell}}^{\circ} = E_{\text{cathode}}^{\circ} - E_{\text{anode}}^{\circ} / E_{\text{sel}}^{\circ} = E_{\text{katode}}^{\circ} - E_{\text{anode}}^{\circ}$	
or/of $E_{\text{cell}}^{\circ} = E_{\text{reduction}}^{\circ} - E_{\text{oxidation}}^{\circ} / E_{\text{sel}}^{\circ} = E_{\text{reduksie}}^{\circ} - E_{\text{oksidasie}}^{\circ}$	
or/of $E_{\text{cell}}^{\circ} = E_{\text{oxidising agent}}^{\circ} - E_{\text{reducing agent}}^{\circ} / E_{\text{sel}}^{\circ} = E_{\text{oksideermiddel}}^{\circ} - E_{\text{reduseermiddel}}^{\circ}$	

TABLE 3: THE PERIODIC TABLE OF ELEMENTS
TABEL 3: DIE PERIODIEKE TABEL VAN ELEMEN

KEY/ISLEUTEL		Atomic number Atoomgetal																	
		2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	
		(I)	(II)	(III)	(IV)	(V)	(VI)	(VII)	(VIII)	(IX)	(X)	(XI)	(XII)	(XIII)	(XIV)	(XV)	(XVI)	(XVII)	(XVIII)
								Electronegativity Elektronegatiwiteit				Approximate relative atomic mass Benaderde relatiewe atoommassa							
		↑		↑				↑				↑				↑			
		←		←				←				←				←			
		↓		↓				↓				↓				↓			
		→		→				→				→				→			
		←		←				←				←				←			
		↓		↓				↓				↓				↓			
		←		←				←				←				←			
		↓		↓				↓				↓				↓			
		←		←				←				←				←			
1	H	1																	
3	Li	7	4																
11	Na	23	24	12															
19	K	39	40	20	Ca	45		21	SC	48									
37	Rb	85	86	38	Sr	88		40	Zr	91									
55	Cs	133	137	56	Ba	137		72	Hf	179									
87	Fr			88	Ra	226		89	AC										
23	Al	27																	
13	Al	27																	
14	Si	28																	
15	P	31																	
16	S	32																	
17	Cl	35,5																	
18	Ar	40																	
23	V	51																	
41	Nb	93																	
73	Ta	181																	
103	Rh	103																	
101	Ru	101																	
106	Pd	106																	
115	In	115																	
81	Tl	204																	
112	Cd	112																	
119	Sb	122																	
127	I	127																	
128	Te	128																	
85	At	209																	
86	Rn	222																	
88	Ra	226																	
90	Th	232																	
91	Pa	231																	
92	U	238																	
93	Np	237																	
94	Pu	244																	
95	Am	243																	
96	Cm	247																	
97	Bk	247																	
98	Cf	251																	
99	Es	252																	
100	Fm	257																	
101	Md	258																	
102	No	259																	
103	Lr	262																	
104	Rf	261																	
105	Db	262																	
106	Sg	266																	
107	Bh	264																	
108	Hs	277																	
109	Mt	268																	
110	Ds	271																	
111	Rg	272																	
112	Cn	285																	
113	Nh	284																	
114	Fl	289																	
115	Mc	288																	
116	Lv	293																	
117	Ts	294																	
118	Og	294																	

Downloaded from Stanmorephysics.com
education



Department:
Education
PROVINCE OF KWAZULU-NATAL

**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

PHYSICAL SCIENCES: CHEMISTRY (P2)

MARKING GUIDELINE

COMMON TEST

MARCH 2019

MARKS: 50

TIME: 1 hour

This marking guideline consists of 5 pages.


QUESTION 1

- 1.1 C ✓✓ (2)
- 1.2 D ✓✓ (2)
- 1.3 A ✓✓ (2)
- 1.4 B ✓✓ (2)
- [8]**



QUESTION 2

- 2.1.1 Hydroxyl group ✓ (1)
- 2.1.2 methylpropan - 1 - ol ✓ Accept 2 – methylpropan - 1 - ol (2)
- 2.1.3 (Any One)

- NOTE:**
- For any additional bond on the carbon atoms subtract 1 mark.
 - If there are missing hydrogens subtract 1 mark.





Downloaded from Stanmorephysics.com



✓✓

(2)

2.2



Reactants ✓ / Products ✓ / Balancing ✓

If structural/condensed formula is used then 2/3 max.

(3)

2.3 $(\frac{12,5}{100}) \times M(C_nH_{2n}O_2) = 32$ ✓

$M(C_nH_{2n}O_2) = 256 \text{ g.mol}^{-1}$ ✓

But $12(n) + 2(n) = 256 - 32$

$n = 16$ ✓

$\therefore \% C = (\frac{16 \times 12}{256}) \times 100$ ✓


$= 75\%$ ✓

(5)

Calc. molar mass of compound ✓✓
Determining the value of n ✓
Calc. % of C in whole comp. ✓✓

2.4

2.4.1



(2)

2.4.2 polyethene ✓✓

(2)

[19]

Downloaded from Stanmorephysics.com

QUESTION 3

3.1 A series of organic compounds that can be described by the same general formula ✓✓

OR

A series of organic compounds in which one member differs from the next by a $-\text{CH}_2$ group. (2)

3.2 What is the relationship between boiling point and type of functional group? ✓✓
(Accept any other correct investigative question as long as there is a relationship between boiling point and functional group.) (2)

3.3 P✓
P has the lowest boiling point. ✓ (2)

3.4 { 3.4.1 $\text{CH}_3(\text{CH}_2)_2\text{COOH}$ ✓ (1)

3.4.2 $\text{CH}_3(\text{CH}_2)_2\text{COOH}$ and $\text{CH}_3(\text{CH}_2)_3\text{CH}_2\text{OH}$ have hydrogen bonding and $\text{CH}_3(\text{CH}_2)\text{CHO}$ has no hydrogen bonding. ✓
 $\text{CH}_3(\text{CH}_2)_2\text{COOH}$ has TWO sites for hydrogen bonding. ✓
The intermolecular forces between molecules of $\text{CH}_3(\text{CH}_2)_2\text{COOH}$ is therefore the strongest ✓ and will require the most amount of energy to separate its molecule. ✓ (4)

3.5 greater than ✓ (1)
[12]

Downloaded from Stanmorephysics.com

QUESTION 4

- 4.1
- 4.1.1 substitution/hydrolysis ✓ (1)
- 4.1.2 dissolve $\text{CH}_3\text{CH}_2\text{CHBrCH}_3$ in ethanol/ethanol as a solvent ✓
mild heat ✓
dilute strong base/NaOH / KOH ✓ (any 2) (2)
- 4.1.3 butan – 2 – ol ✓✓
Accept 2-butanol (2)
- 4.2
- 4.2.1 dehydration ✓ (1)
- 4.2.2 water ✓ (1)
- 4.2.3 major product ✓ (1)
- 4.3
- 4.3.1 addition/hydrohalogenation ✓ (1)
- 4.3.2 Prevents the formation of an alcohol ✓✓ (2)
- [11]

TOTAL MARKS: [50]

Downloaded from Stanmorephysics.com