



education

DEPARTMENT: EDUCATION
MPUMALANGA PROVINCE

**NATIONAL
SENIOR CERTIFICATE**

GRADE 12

PHYSICAL SCIENCES: CHEMISTRY P2

SEPTEMBER 2019

MARKS: 150

TIME: 3 hour

This question paper consists of 15 pages and 4 data sheets.

INSTRUCTIONS AND INFORMATION

1. Write your name in the appropriate space on the ANSWER BOOK.
2. This question paper consists of TEN questions. Answer ALL the questions in the ANSWER BOOK.
3. Start EACH question on a NEW page in the ANSWER BOOK.
4. Number the answers correctly according to the numbering system used in this question paper.
5. Leave ONE line between two sub questions, for example between QUESTION 2.1 and QUESTION 2.2.
6. You may use a non-programmable calculator.
7. You may use appropriate mathematical instruments.
8. You are advised to use the attached DATA SHEETS.
9. Show ALL formulae and substitutions in ALL calculations.
10. Round off your final numerical answers to a MINIMUM of TWO decimal places.
11. Give brief motivations, discussions, et cetera where required.
12. Write neatly and legibly.

QUESTION 1 : MULTIPLE-CHOICE QUESTIONS

Four options are provided as possible answers to the following questions. Each question has only ONE correct answer. Choose the answer and write only the letter (A – D) next to the question number (1.1 – 1.10) in the ANSWER BOOK , for example, 1.11 E.

1.1 Consider the following statements regarding organic compounds belonging to the same homologous series.

- I They have the same boiling point.
- II They have the same functional group.
- III They have the same molecular formula.

Which of the above statements is/are TRUE?

- A II only
- B I and II only
- C II and III only
- D I, II and III

(2)

1.2 Which ONE of the following organic compounds is a haloalkane?

- A Methane
- B Tetrachloromethane
- C Methyl methanoate
- D Methanal

(2)

- 1.3 The organic compounds shown in the table below have the same molar mass ($60 \text{ g}\cdot\text{mol}^{-1}$).


| Compound | Structural formula | Boiling point ($^{\circ}\text{C}$) |
|----------|---|--------------------------------------|
| X | $ \begin{array}{ccccccc} & \text{H} & & \text{H} & & \text{H} & \\ & & & & & & \\ \text{H} & -\text{C} & - & \text{C} & - & \text{C} & -\text{O}-\text{H} \\ & & & & & & \\ & \text{H} & & \text{H} & & \text{H} & \end{array} $ | 97 |
| Y | $ \begin{array}{ccccccc} & & & \text{O} & & & \\ & & & & & & \\ & \text{H} & & & & & \\ & & & & & & \\ \text{H} & -\text{C} & - & \text{C} & -\text{O}-\text{H} \\ & & & & & & \\ & \text{H} & & & & & \end{array} $ | ? |
| Z | $ \begin{array}{ccccccc} & & & \text{O} & & \text{H} & \\ & & & & & & \\ \text{H} & -\text{C} & -\text{O}- & \text{C} & -\text{H} \\ & & & & & & \\ & & & \text{H} & & \text{H} & \end{array} $ | 32 |

Which ONE of the following is most likely to be the boiling point (in $^{\circ}\text{C}$) of compound Y?

- A 24
B 40
C 90
D 118

(2)

- 1.4 The reaction represented below reaches equilibrium in a closed container.



Which ONE of the following changes in equilibrium conditions will result in an INCREASE in the value of the equilibrium constant (K_c)?

- A Decreasing the pressure by increasing the volume
B Decreasing the temperature
C Increasing the pressure by decreasing the volume
D Increasing the temperature

(2)

- 1.5 Four different solutions, each with concentration $0,1 \text{ mol}\cdot\text{dm}^{-3}$, are prepared. Which ONE of the solutions will have the lowest pH?

- A NH_4NO_3
 B Na_2CO_3
 C KNO_3
 D KOH


(2)

- 1.6 A table of acid-base indicators and the pH ranges over which they change colour are given below. Which ONE of these indicators is most suitable for use in the titration of ethanoic acid with sodium hydroxide?

| | Indicator | pH range |
|---|------------------|-----------|
| A | Bromophenol blue | 3,0 – 4,6 |
| B | Litmus | 4,5 – 8,3 |
| C | Bromothymol blue | 6,0 – 7,6 |
| D | Cresolphthalein | 8,2 – 9,8 |

(2)

- 1.7 Two hypothetical half-reactions and their respective standard reduction potentials are given below.



Which ONE of the following has the greatest tendency to donate electrons?

- A X^-
 B X_2
 C Y
 D Y^+


(2)

- 1.8 Which ONE of the following cell reactions will be spontaneous?

- A $\text{Sn}^{2+}(\text{aq}) + \text{Zn}^{2+}(\text{aq}) \rightarrow \text{Zn}(\text{s}) + \text{Sn}^{4+}(\text{aq})$
 B $2\text{H}^+(\text{aq}) + \text{Cu}(\text{s}) \rightarrow \text{Cu}^{2+}(\text{aq}) + \text{H}_2(\text{g})$
 C $\text{Cd}(\text{s}) + 2\text{Co}^{3+}(\text{aq}) \rightarrow \text{Cd}^{2+}(\text{aq}) + 2\text{Co}^{2+}(\text{aq})$
 D $2\text{Cl}^-(\text{aq}) + 2\text{Fe}^{3+}(\text{aq}) \rightarrow \text{Cl}_2(\text{g}) + 2\text{Fe}^{2+}(\text{aq})$

(2)

1.9 The diagram below represents a cell used to electroplate a key with silver metal.



Which ONE of the following combinations represent the CHANGE in mass of the silver electrode and the concentration of the AgNO_3 electrolyte during electrolysis?

| | Mass of silver electrode | Concentration of AgNO_3 (aq) |
|---|--------------------------|---------------------------------------|
| A | Decreases | Increases |
| B | Decreases | No change |
| C | Increases | No change |
| D | Increases | Increases |

(2)


1.10 Which combination of fertilisers will best address the following needs of a crop?

| | Enhance root growth | Assist with ripening of fruit |
|---|--|--|
| A | K_2SO_4 | H_2NCONH_2 |
| B | KCl | $\text{Ca}_3(\text{H}_2\text{PO}_4)_2$ |
| C | $\text{Ca}_3(\text{H}_2\text{PO}_4)_2$ | KCl |
| D | NH_4NO_3 | $\text{Ca}_3(\text{H}_2\text{PO}_4)_2$ |

(2)
[20]

QUESTION 2 (Start on a new page.)

2.1 The structural formula of an unsaturated hydrocarbon is given below.



2.1.1 Define the term *unsaturated hydrocarbon*. (2)

2.1.2 Write down the IUPAC name of the above compound. (2)

2.1.3 Write down the IUPAC name of a positional isomer of the above compound. (2)

The hydrocarbon above reacts with bromine water.

2.1.4 Use structural formulae and write down a balanced equation for the reaction. (3)

2.1.5 Write down the name of the type of reaction that takes place. (1)

2.1.6 What colour change will be observed during this reaction? (1)

2.2 Butanoic acid is a functional isomer of methyl propanoate.

2.2.1 Define the term *functional isomer*. (2)

2.2.2 Draw the structural formula of butanoic acid. (2)

2.2.3 Draw the structural formula of methyl propanoate. (2)

2.2.4 Write down the IUPAC name of the alcohol used to prepare methyl propanoate. (1)

2.2.5 Write down TWO reaction conditions needed for the preparation of an ester. (2)


2.2.6 Which one of BUTANOIC ACID or METHYL PROPANOATE will have the higher boiling point? (1)

2.2.7 Explain the answer in QUESTION 2.2.6. (4)

[25]

QUESTION 3 (Start on a new page.)

Consider the organic reactions (A to F) represented in the flow diagram below.



- 3.1 Write down the type of reaction represented by:
- 3.1.1 **A** (1)
- 3.1.2 **C** (1)
- 3.2 Using structural formulae, write down the balanced equation for reaction **D**. (4)
- 3.3 Write down the IUPAC name of compound **X**. (2)
- 3.4 Write down the NAME or FORMULA of the reagent that is needed in reaction **F**. (1)
- 3.5 What TYPE of addition reaction is represented by reaction **E**? (1)
- 3.6 Using MOLECULAR formulae, write down the balanced equation for the complete combustion of the saturated hydrocarbon formed in reaction **E**. (4)
- [14]

QUESTION 4 (Start on a new page.)


The boiling points of four organic compounds are given in the table below.

| | Compound | Boiling point (°C) |
|----------|-------------|--------------------|
| A | Ethanol | 79 |
| B | Propan-1-ol | 98 |
| C | Butan-1-ol | 117 |
| D | Pentan-1-ol | 138 |

- 4.1 Write down the NAME of the functional group of the compounds **A** to **D**. (1)
- 4.2 Is butan-1-ol a PRIMARY or a SECONDARY alcohol? (1)
- 4.3 Define the term *boiling point*. (2)
- 4.4 Use the table to explain the difference in boiling points of compounds **A** to **D**. (3)
- 4.5 Which one of compounds, **A** or **B**, has the higher vapour pressure? Give a reason the answer. (2)
- [9]

QUESTION 5 (Start on a new page.)

Learners use the reaction of $\text{Na}_2\text{S}_2\text{O}_3(\text{aq})$ and $\text{HCl}(\text{aq})$ to investigate different factors which influence reaction rate. The balanced equation for the reaction is:



Equal volumes of the two solutions are mixed and the time taken for a certain amount of sulphur to form is measured.

5.1 The reaction conditions used and the results obtained during the first investigation are recorded in the table below.

| | Temperature (°C) | Concentration of $\text{Na}_2\text{S}_2\text{O}_3$ ($\text{mol}\cdot\text{dm}^{-3}$) | Concentration of HCl ($\text{mol}\cdot\text{dm}^{-3}$) | Time (s) |
|---------------------|------------------|--|---|----------|
| Experiment 1 | 20 | 0,5 | 0,5 | 40 |
| Experiment 2 | 20 | 0,9 | 0,5 | 25 |
| Experiment 3 | 20 | 1,4 | 0,5 | 15 |

Use the information in the table to answer the following questions.
For this investigation, write down:

- 5.1.1 One controlled variable (1)
- 5.1.2 The independent variable (1)
- 5.1.3 The conclusion that can be drawn from the results (2)
- 5.1.4 The limiting reactant ($\text{Na}_2\text{S}_2\text{O}_3$ or HCl) in **Experiment 1**. Explain the answer. (4)


5.2 The reaction conditions used and the results obtained during a second investigation are recorded in the table below.

| | Temperature (°C) | Concentration of $\text{Na}_2\text{S}_2\text{O}_3$ ($\text{mol}\cdot\text{dm}^{-3}$) | Concentration of HCl ($\text{mol}\cdot\text{dm}^{-3}$) | Time (s) |
|---------------------|------------------|--|---|----------|
| Experiment 4 | 20 | 0,5 | 0,5 | 40 |
| Experiment 5 | 30 | 0,5 | 0,5 | 20 |
| Experiment 6 | 40 | 0,5 | 0,5 | 10 |

- 5.2.1 Which ONE of the three experiments (4, 5 or 6) has the highest rate of reaction? (1)
- 5.2.2 Explain the answer in QUESTION 5.2.1 in terms of the collision theory. (3)
- [12]**

QUESTION 6 (Start on a new page.)


- 6.1 Potassium chromate (K_2CrO_4) is dissolved in water. In solution, the chromate ions (CrO_4^{2-}) reach equilibrium with the dichromate ions ($Cr_2O_7^{2-}$) according to the following balanced equation:




- 6.1.1 A reagent is now added to the orange solution. Which reagent, $NaOH(aq)$ or $HNO_3(aq)$, will result in a colour change from orange to yellow? (1)
- 6.1.2 Which ion, $CrO_4^{2-}(aq)$ or $Cr_2O_7^{2-}(aq)$, will be present in higher concentration at a low pH? (1)

When the temperature is gradually increased, it is observed that the colour of the solution changes to yellow.

- 6.1.3 Is the forward reaction ENDOTHERMIC or EXOTHERMIC? (1)
- 6.1.4 Explain the answer in QUESTION 6.1.3 by referring to Le Chatelier's principle. (2)
- 6.2 5 moles of hydrogen gas, $H_2(g)$, and 5 moles of iodine vapour, $I_2(g)$, are sealed in a 2 dm^3 vessel at a temperature of 600 K. The reaction reaches equilibrium according to the following balanced equation:



- 6.2.1 Calculate the concentration of the hydrogen iodide (HI) at equilibrium if the equilibrium constant, K_c , for the reaction at 600 K is 0,36. (8)
- 6.2.2 The pressure on the system is now changed. The graph below shows the change in concentrations of the reactants and products as a result of this change.





Was the pressure INCREASED or DECREASED at t_1 ? Explain the answer.


(3)
[16]

QUESTION 7 (Start on a new page.)

- 7.1 Reactions **A** and **B** below represent the ionisation of two acids, HF(aq) and HBr(aq) respectively. The ionisation constants (K_a) for the two acids are given at 25 °C.



- 7.1.1 Write down the NAME or FORMULA of the conjugate base of HF(aq). (1)
- 7.1.2 Define the term *weak acid*. (2)
- 7.1.3 Which acid, HF(aq) or HBr(aq), is stronger? Explain your choice. (3)
- 7.1.4 Water can act as an ampholyte. Define the term *ampholyte*. (2)
- 7.2 7,5 g IMPURE sodium hydroxide pellets, NaOH(s), is added to a flask containing 700 cm³ of a 0,15 mol·dm⁻³ sulphuric acid solution, H₂SO₄(aq). The balanced equation for the reaction is:



- 7.2.1 Calculate the number of moles of sulphuric acid in the flask. (3)
- 7.2.2 The percentage purity of the sodium hydroxide pellets used is 80%. Calculate the pH of the solution in the flask after the addition of the NaOH(s). (7)


[18]

QUESTION 8 (Start on a new page.)

- 8.1 An iron nail, which has been thoroughly cleaned, is placed in a beaker containing a $1 \text{ mol}\cdot\text{dm}^{-3}$ copper(II) sulphate solution. The beaker is allowed to stand overnight. The next morning it is observed that a reddish brown precipitate formed on the iron nail.

By referring to the relative strength of reducing agents, explain why a reddish brown precipitate formed on the iron nail. (3)

- 8.2 A galvanic cell is set up under standard conditions as shown in the simplified diagram below.




- 8.2.1 State ONE function of the component labelled Q. (1)
- 8.2.2 Which electrode (Cu or Cr) is the cathode of this cell? (1)
- 8.2.3 Write down the overall (net) balanced cell reaction that takes place in this cell. (3)
- 8.2.4 Calculate the initial emf of this cell. (4)
- 8.2.5 A similar cell is now set up using $2 \text{ mol}\cdot\text{dm}^{-3} \text{ Cu}^{2+}(\text{aq})$. All other conditions stay the same. How will the initial emf of this cell compare to that calculated in QUESTION 8.2.4? Choose from HIGHER THAN, LOWER THAN or EQUAL TO. Explain the answer. (3)
- 8.2.6 How will the initial reading on the voltmeter change as the cell reaction approaches equilibrium? Choose from INCREASES, DECREASES or REMAINS THE SAME. (1)

[16]

QUESTION 9 (Start on a new page.)

A simplified diagram of an electrochemical cell, used for the production of chlorine gas, is shown below.




- 9.1 Define the term *electrolyte*. (2)
- 9.2 Write down the NAME of electrolyte X. (1)
- 9.3 Write down the half-reaction which takes place at the positive electrode. (2)
- 9.4 Which electrode, **A** or **B**, is the ANODE? Give a reason for the answer. (2)
- 9.5 Litmus added to the electrolyte turns blue around electrode **B**. Explain this observation with the aid of the relevant half-reaction. (3)
- 9.6 Give a reason why platinum can be used as electrodes. (1)

[11]

QUESTION 10 (Start on a new page.)

Two processes in the fertiliser industry are summarised as in the flow diagram below.



10.1 Use the information above and write down:

10.1.1 NAME of **Process 1**

(1)

10.1.2 NAME or FORMULA of compound **Y**

(1)

10.1.3 Write down a balanced chemical equation for the preparation of fertiliser **Z**.

(3)

10.2 Calculate the number of moles of nitrogen in a 5 kg bag of fertiliser with a ratio 7:1:3(22).

(4)

[9]

TOTAL: 150

**DATA FOR PHYSICAL SCIENCES GRADE 12
PAPER 2 (CHEMISTRY)**

**GEGEWENS VIR FISIESTE WETENSAPPE GRAAD 12
VRAESTEL 2 (CHEMIE)**

TABLE 1: PHYSICAL CONSTANTS/TABEL 1: FISIESTE KONSTANTES

| NAME/NAAM | SYMBOL/SIMBOOL | VALUE/WAARDE |
|--|----------------|---|
| Standard pressure Standaarddruk | p^{θ} | $1,013 \times 10^5 \text{ Pa}$ |
| Molar gas volume at STP Molêre gasvolume by STD | V_m | $22,4 \text{ dm}^3 \cdot \text{mol}^{-1}$ |
| Standard temperature Standaardtemperatuur | T^{θ} | 273 K |
| Avogadro's constant | N_A | $6,023 \times 10^{23} \text{ mol}^{-1}$ |

TABLE 2: FORMULAE/TABEL 2: FORMULES

| | |
|---|---|
| $n = \frac{m}{M}$ | $n = \frac{N}{N_A}$ |
| $c = \frac{n}{V}$ OR/OF $c = \frac{m}{MV}$ | $n = \frac{V}{V_m}$ |
| $\frac{n_a}{n_b} = \frac{c_a V_a}{c_b V_b}$ | $\text{pH} = -\log[\text{H}_3\text{O}^+]$ |
| $K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1 \times 10^{-14}$ at/by 298 K | |
| $E_{\text{cell}}^{\theta} = E_{\text{cathode}}^{\theta} - E_{\text{anode}}^{\theta}$ / $E_{\text{sel}}^{\theta} = E_{\text{katode}}^{\theta} - E_{\text{anode}}^{\theta}$ | |
| OR/OF | |
| $E_{\text{cell}}^{\theta} = E_{\text{reduction}}^{\theta} - E_{\text{oxidation}}^{\theta}$ / $E_{\text{sel}}^{\theta} = E_{\text{reduksie}}^{\theta} - E_{\text{oksidasie}}^{\theta}$ | |
| OR/OF | |
| $E_{\text{cell}}^{\theta} = E_{\text{oxidisingagent}}^{\theta} - E_{\text{reducingagent}}^{\theta}$ / $E_{\text{sel}}^{\theta} = E_{\text{oksideermiddel}}^{\theta} - E_{\text{reduseermiddel}}^{\theta}$ | |

TABLE 4A: STANDARD REDUCTION POTENTIALS
TABEL 4A: STANDAARD-REDUKSIEPOTENSIALE

| Half-reactions/ <i>Halfreaksies</i> | E^θ (V) |
|---|----------------|
| $F_2(g) + 2e^- \rightleftharpoons 2F^-$ | + 2,87 |
| $Co^{3+} + e^- \rightleftharpoons Co^{2+}$ | + 1,81 |
| $H_2O_2 + 2H^+ + 2e^- \rightleftharpoons 2H_2O$ | + 1,77 |
| $MnO_4^- + 8H^+ + 5e^- \rightleftharpoons Mn^{2+} + 4H_2O$ | + 1,51 |
| $Cl_2(g) + 2e^- \rightleftharpoons 2Cl^-$ | + 1,36 |
| $Cr_2O_7^{2-} + 14H^+ + 6e^- \rightleftharpoons 2Cr^{3+} + 7H_2O$ | + 1,33 |
| $O_2(g) + 4H^+ + 4e^- \rightleftharpoons 2H_2O$ | + 1,23 |
| $MnO_2 + 4H^+ + 2e^- \rightleftharpoons Mn^{2+} + 2H_2O$ | + 1,23 |
| $Pt^{2+} + 2e^- \rightleftharpoons Pt$ | + 1,20 |
| $Br_2(l) + 2e^- \rightleftharpoons 2Br^-$ | + 1,07 |
| $NO_3^- + 4H^+ + 3e^- \rightleftharpoons NO(g) + 2H_2O$ | + 0,96 |
| $Hg^{2+} + 2e^- \rightleftharpoons Hg(l)$ | + 0,85 |
| $Ag^+ + e^- \rightleftharpoons Ag$ | + 0,80 |
| $NO_3^- + 2H^+ + e^- \rightleftharpoons NO_2(g) + H_2O$ | + 0,80 |
| $Fe^{3+} + e^- \rightleftharpoons Fe^{2+}$ | + 0,77 |
| $O_2(g) + 2H^+ + 2e^- \rightleftharpoons H_2O_2$ | + 0,68 |
| $I_2 + 2e^- \rightleftharpoons 2I^-$ | + 0,54 |
| $Cu^+ + e^- \rightleftharpoons Cu$ | + 0,52 |
| $SO_2 + 4H^+ + 4e^- \rightleftharpoons S + 2H_2O$ | + 0,45 |
| $2H_2O + O_2 + 4e^- \rightleftharpoons 4OH^-$ | + 0,40 |
| $Cu^{2+} + 2e^- \rightleftharpoons Cu$ | + 0,34 |
| $SO_4^{2-} + 4H^+ + 2e^- \rightleftharpoons SO_2(g) + 2H_2O$ | + 0,17 |
| $Cu^{2+} + e^- \rightleftharpoons Cu^+$ | + 0,16 |
| $Sn^{4+} + 2e^- \rightleftharpoons Sn^{2+}$ | + 0,15 |
| $S + 2H^+ + 2e^- \rightleftharpoons H_2S(g)$ | + 0,14 |
| $2H^+ + 2e^- \rightleftharpoons H_2(g)$ | 0,00 |
| $Fe^{3+} + 3e^- \rightleftharpoons Fe$ | - 0,06 |
| $Pb^{2+} + 2e^- \rightleftharpoons Pb$ | - 0,13 |
| $Sn^{2+} + 2e^- \rightleftharpoons Sn$ | - 0,14 |
| $Ni^{2+} + 2e^- \rightleftharpoons Ni$ | - 0,27 |
| $Co^{2+} + 2e^- \rightleftharpoons Co$ | - 0,28 |
| $Cd^{2+} + 2e^- \rightleftharpoons Cd$ | - 0,40 |
| $Cr^{3+} + e^- \rightleftharpoons Cr^{2+}$ | - 0,41 |
| $Fe^{2+} + 2e^- \rightleftharpoons Fe$ | - 0,44 |
| $Cr^{3+} + 3e^- \rightleftharpoons Cr$ | - 0,74 |
| $Zn^{2+} + 2e^- \rightleftharpoons Zn$ | - 0,76 |
| $2H_2O + 2e^- \rightleftharpoons H_2(g) + 2OH^-$ | - 0,83 |
| $Cr^{2+} + 2e^- \rightleftharpoons Cr$ | - 0,91 |
| $Mn^{2+} + 2e^- \rightleftharpoons Mn$ | - 1,18 |
| $Al^{3+} + 3e^- \rightleftharpoons Al$ | - 1,66 |
| $Mg^{2+} + 2e^- \rightleftharpoons Mg$ | - 2,36 |
| $Na^+ + e^- \rightleftharpoons Na$ | - 2,71 |
| $Ca^{2+} + 2e^- \rightleftharpoons Ca$ | - 2,87 |
| $Sr^{2+} + 2e^- \rightleftharpoons Sr$ | - 2,89 |
| $Ba^{2+} + 2e^- \rightleftharpoons Ba$ | - 2,90 |
| $Cs^+ + e^- \rightleftharpoons Cs$ | - 2,92 |
| $K^+ + e^- \rightleftharpoons K$ | - 2,93 |
| $Li^+ + e^- \rightleftharpoons Li$ | - 3,05 |

Increasing oxidising ability/*Toenemende oksiderende vermoë*

Increasing reducing ability/*Toenemende reduserende vermoë*

TABLE 4B: STANDARD REDUCTION POTENTIALS
TABEL 4B: STANDAARD-REDUKSIEPOTENSIALE

| Half-reactions/ <i>Halfreaksies</i> | E^{θ} (V) |
|--|---------------------|
| $\text{Li}^+ + \text{e}^- \rightleftharpoons \text{Li}$ | -3,05 |
| $\text{K}^+ + \text{e}^- \rightleftharpoons \text{K}$ | -2,93 |
| $\text{Cs}^+ + \text{e}^- \rightleftharpoons \text{Cs}$ | -2,92 |
| $\text{Ba}^{2+} + 2\text{e}^- \rightleftharpoons \text{Ba}$ | -2,90 |
| $\text{Sr}^{2+} + 2\text{e}^- \rightleftharpoons \text{Sr}$ | -2,89 |
| $\text{Ca}^{2+} + 2\text{e}^- \rightleftharpoons \text{Ca}$ | -2,87 |
| $\text{Na}^+ + \text{e}^- \rightleftharpoons \text{Na}$ | -2,71 |
| $\text{Mg}^{2+} + 2\text{e}^- \rightleftharpoons \text{Mg}$ | -2,36 |
| $\text{Al}^{3+} + 3\text{e}^- \rightleftharpoons \text{Al}$ | -1,66 |
| $\text{Mn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Mn}$ | -1,18 |
| $\text{Cr}^{2+} + 2\text{e}^- \rightleftharpoons \text{Cr}$ | -0,91 |
| $2\text{H}_2\text{O} + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g}) + 2\text{OH}^-$ | -0,83 |
| $\text{Zn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Zn}$ | -0,76 |
| $\text{Cr}^{3+} + 3\text{e}^- \rightleftharpoons \text{Cr}$ | -0,74 |
| $\text{Fe}^{2+} + 2\text{e}^- \rightleftharpoons \text{Fe}$ | -0,44 |
| $\text{Cr}^{3+} + \text{e}^- \rightleftharpoons \text{Cr}^{2+}$ | -0,41 |
| $\text{Cd}^{2+} + 2\text{e}^- \rightleftharpoons \text{Cd}$ | -0,40 |
| $\text{Co}^{2+} + 2\text{e}^- \rightleftharpoons \text{Co}$ | -0,28 |
| $\text{Ni}^{2+} + 2\text{e}^- \rightleftharpoons \text{Ni}$ | -0,27 |
| $\text{Sn}^{2+} + 2\text{e}^- \rightleftharpoons \text{Sn}$ | -0,14 |
| $\text{Pb}^{2+} + 2\text{e}^- \rightleftharpoons \text{Pb}$ | -0,13 |
| $\text{Fe}^{3+} + 3\text{e}^- \rightleftharpoons \text{Fe}$ | -0,06 |
| $2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g})$ | 0,00 |
| $\text{S} + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2\text{S}(\text{g})$ | +0,14 |
| $\text{Sn}^{4+} + 2\text{e}^- \rightleftharpoons \text{Sn}^{2+}$ | +0,15 |
| $\text{Cu}^{2+} + \text{e}^- \rightleftharpoons \text{Cu}^+$ | +0,16 |
| $\text{SO}_4^{2-} + 4\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{SO}_2(\text{g}) + 2\text{H}_2\text{O}$ | +0,17 |
| $\text{Cu}^{2+} + 2\text{e}^- \rightleftharpoons \text{Cu}$ | +0,34 |
| $2\text{H}_2\text{O} + \text{O}_2 + 4\text{e}^- \rightleftharpoons 4\text{OH}^-$ | +0,40 |
| $\text{SO}_2 + 4\text{H}^+ + 4\text{e}^- \rightleftharpoons \text{S} + 2\text{H}_2\text{O}$ | +0,45 |
| $\text{Cu}^+ + \text{e}^- \rightleftharpoons \text{Cu}$ | +0,52 |
| $\text{I}_2 + 2\text{e}^- \rightleftharpoons 2\text{I}^-$ | +0,54 |
| $\text{O}_2(\text{g}) + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{H}_2\text{O}_2$ | +0,68 |
| $\text{Fe}^{3+} + \text{e}^- \rightleftharpoons \text{Fe}^{2+}$ | +0,77 |
| $\text{NO}_3^- + 2\text{H}^+ + \text{e}^- \rightleftharpoons \text{NO}_2(\text{g}) + \text{H}_2\text{O}$ | +0,80 |
| $\text{Ag}^+ + \text{e}^- \rightleftharpoons \text{Ag}$ | +0,80 |
| $\text{Hg}^{2+} + 2\text{e}^- \rightleftharpoons \text{Hg}(\text{l})$ | +0,85 |
| $\text{NO}_3^- + 4\text{H}^+ + 3\text{e}^- \rightleftharpoons \text{NO}(\text{g}) + 2\text{H}_2\text{O}$ | +0,96 |
| $\text{Br}_2(\text{l}) + 2\text{e}^- \rightleftharpoons 2\text{Br}^-$ | +1,07 |
| $\text{Pt}^{2+} + 2\text{e}^- \rightleftharpoons \text{Pt}$ | +1,20 |
| $\text{MnO}_2 + 4\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{Mn}^{2+} + 2\text{H}_2\text{O}$ | +1,23 |
| $\text{O}_2(\text{g}) + 4\text{H}^+ + 4\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}$ | +1,23 |
| $\text{Cr}_2\text{O}_7^{2-} + 14\text{H}^+ + 6\text{e}^- \rightleftharpoons 2\text{Cr}^{3+} + 7\text{H}_2\text{O}$ | +1,33 |
| $\text{Cl}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{Cl}^-$ | +1,36 |
| $\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^- \rightleftharpoons \text{Mn}^{2+} + 4\text{H}_2\text{O}$ | +1,51 |
| $\text{H}_2\text{O}_2 + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}$ | +1,77 |
| $\text{Co}^{3+} + \text{e}^- \rightleftharpoons \text{Co}^{2+}$ | +1,81 |
| $\text{F}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{F}^-$ | +2,87 |

Increasing oxidising ability/*Toenemende oksiderende vermoë*Increasing reducing ability/*Toenemende reduserende vermoë*

TABLE 3: THE PERIODIC TABLE OF ELEMENTS
TABLE 3: THE PERIODIC TABLE OF ELEMENTS

| | | KEY/SLEUTEL | | | | | | | | | | | | | | | | | |
|-----------------|-----------------|--|-----------------|-----------------|----------------|-----------------|-----------------|-----------------|-----------------|------------------|-----------------|-----------------|-----------------|-----------------|-----------------|------------------|-----------------|--|--|
| | | Atomic number Atoomgetal | | | | | | | | | | | | | | | | | |
| | | Electronegativity Elektronegatiwiteit | | | | | | | | | | | | | | | | | |
| | | Approximate relative atomic mass Benaderde relatiewe atoommassa | | | | | | | | | | | | | | | | | |
| | | Symbol Simbool | | | | | | | | | | | | | | | | | |
| (I) | (II) | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 (III) | 14 (IV) | 15 (V) | 16 (VI) | 17 (VII) | 18 (VIII) | | |
| 1 H 1 | 2 He 4 | 3 Li 7 | 4 Be 9 | 5 B 11 | 6 C 12 | 7 N 14 | 8 O 16 | 9 F 19 | 10 Ne 20 | 11 Na 23 | 12 Mg 24 | 13 Al 27 | 14 Si 28 | 15 P 31 | 16 S 32 | 17 Cl 35,5 | 18 Ar 40 | | |
| 19 K 39 | 20 Ca 40 | 21 Sc 45 | 22 Ti 48 | 23 V 51 | 24 Cr 52 | 25 Mn 55 | 26 Fe 56 | 27 Co 59 | 28 Ni 59 | 29 Cu 63,5 | 30 Zn 65 | 31 Ga 70 | 32 Ge 73 | 33 As 75 | 34 Se 79 | 35 Br 80 | 36 Kr 84 | | |
| 37 Rb 86 | 38 Sr 88 | 39 Y 89 | 40 Zr 91 | 41 Nb 92 | 42 Mo 96 | 43 Tc 98 | 44 Ru 101 | 45 Rh 103 | 46 Pd 106 | 47 Ag 108 | 48 Cd 112 | 49 In 115 | 50 Sn 119 | 51 Sb 122 | 52 Te 128 | 53 I 127 | 54 Xe 131 | | |
| 55 Cs 133 | 56 Ba 137 | 57 La 139 | 72 Hf 179 | 73 Ta 181 | 74 W 184 | 75 Re 186 | 76 Os 190 | 77 Ir 192 | 78 Pt 195 | 79 Au 197 | 80 Hg 201 | 81 Tl 204 | 82 Pb 207 | 83 Bi 209 | 84 Po 209 | 85 At 210 | 86 Rn 222 | | |
| 87 Fr 223 | 88 Ra 226 | 89 Ac | | | | | | | | | | | | | | | | | |

| | | | | | | | | | | | | | |
|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|------------------|------------------|------------------|------------------|
| 58 Ce 140 | 59 Pr 141 | 60 Nd 144 | 61 Pm | 62 Sm 150 | 63 Eu 152 | 64 Gd 157 | 65 Tb 159 | 66 Dy 163 | 67 Ho 165 | 68 Er 167 | 69 Tm 169 | 70 Yb 173 | 71 Lu 175 |
| 90 Th 232 | 91 Pa 231 | 92 U 238 | 93 Np 237 | 94 Pu 244 | 95 Am 243 | 96 Cm 247 | 97 Bk 247 | 98 Cf 251 | 99 Es 252 | 100 Fm 257 | 101 Md 288 | 102 No 289 | 103 Lr 260 |

