



Province of the  
**EASTERN CAPE**  
EDUCATION

**NATIONAL SENIOR  
CERTIFICATE/  
NASIONALE SENIOR  
SERTIFIKAAT**

**GRADE/GRAAD 12**

**JUNE/JUNIE 2018**

**PHYSICAL SCIENCES P2/FISIESE WETENSKAPPE V2  
MARKING GUIDELINE/NASIENRIGLYN**

**MARKS/PUNTE: 150**

---

This marking guideline consists of 10 pages./ *Hierdie nasienriglyn bestaan uit 10 bladsye.*

---

### QUESTION/VRAAG 1

- 1.1 D ✓✓ (2)
- 1.2 C ✓✓ (2)
- 1.3 B ✓✓ (2)
- 1.4 D ✓✓ (2)
- 1.5 B ✓✓ (2)
- 1.6 D ✓✓ (2)
- 1.7 A ✓✓ (2)
- 1.8 A ✓✓ (2)
- 1.9 B ✓✓ (2)
- 1.10 D ✓✓ (2)
- [20]**

### QUESTION/VRAAG 2


- 2.1.1 Organic compound that consist of carbon and hydrogen atoms only. ✓✓  
*Organiese verbindings wat slegs uit koolstof- en waterstofatome bestaan.*  
**(2 or/of 0)** (2)
- 2.1.2 Alkanes ✓/*Alkane* (1)
- 2.1.3 C<sub>2</sub>H<sub>5</sub> ✓ (1)
- 2.1.4  $2C_4H_{10} + 13O_2 \checkmark \rightarrow 8CO_2 + 10H_2O \checkmark$  Bal ✓

**Notes/Aantekeninge:**

- Reactants✓ Products✓ Balancing✓  
*Reaktanse Produkte Balansering*
- Ignore double arrows and phases./ *Ignoreer dubbelpyle en fases.*
- Marking rule 6.3.10/*Nasienreël 6.3.10.*
- If condensed structural formulae used:/*Indien gekondenseerde struktuurformules gebruik:* Max./Maks. 2/3

- 2.1.5 EXOTHERMIC ✓/*EKSOTERMIES* (1)
- 2.1.6 The chemical process in which longer chain hydrocarbon molecules are broken✓ into shorter more useful molecules. ✓  
*Die chemiese proses waarin langer ketting koolwaterstofmolekules afgebreek word in korter meer bruikbare molekules.* (2)

2.1.7



**Marking criteria/Nasienriglyne:**

- Whole structure correct/ *Hele struktuur korrek*: 2/2
- Only functional group correct.  
*Slegs funksionele groep korrek.* Max./Maks.. 1/2

(2)

2.1.8 CATALYTIC✓//KATALITIES

(1)

2.1.9 UNSATURATED✓//ONVERSADIG

Contains double bonds OR multiple bonds between C atoms.✓  
Bevat dubbelbindings OF meervoudige bindings tussen C-atome.

(2)


2.1.10 2,3-dimethylbut-2-ene/2,3-dimethyl-2-buteen  
2,3-dimetielbut-2-een/2,3-dimetiel-2-buteen

**Marking Criteria/Nasienriglyne:**

- Correct stem i.e but-2-ene/2-butene.  
*Korrekte stam bv.but-2-een/2-buteen.* ✓
- Substituent dimethyl correctly identified. ✓  
*Substituent dimetiel korrek geïdentifiseer.*
- Substituents correctly numbered, hyphens and commas correctly used. ✓  
*Substituente korrek genommer, koppeltekens en kommas korrek gebruik.*

(3)

2.2.1



**Marking criteria/Nasienriglyne:**

- Functional group/*Funksionele groep.* ✓
- Methyl and ethyl substituents /  
*Metiel en etiel substituate* ✓
- Whole structure correct/  
*Hele struktuur korrek* ✓

3/3

(3)

2.2.2 Pentan-2-one✓✓/2-pentanone  
*Pentan-2-oon/2-pentanoon*

(2)

[23]

### QUESTION/VRAAG 3

3.1 A bond or an atom or a group of atoms that determine(s) the physical and chemical properties of a group of organic compounds. ✓✓  
*’n Binding of ’n atoom of ’n groep atome wat die fisiese en chemiese eienskappe van ’n groep organiese verbindings bepaal. (2 or/of 0) (2)*

3.2.1 Aldehyde ✓/Aldehyd (1)

3.2.2 Carboxile group ✓/Karboksiel-groep (1)

3.3.1 185,4(°C) ✓ (1)

 3.3.2

- Compound **B**/carboxylic acid has hydrogen bonding ✓ (in addition to London forces/Dispersion forces/Induced dipole forces/dipole-dipole forces.).  
*Verbinding **B**/karboksielsuur het waterstofbindings behalwe Londonkragte/Dispersiekragte/Geïnduseerde dipoolkragte en dipool-dipoolkragte)*
- Hydrogen bonds are stronger ✓ than London forces/Dispersion forces/Induced dipole forces and dipole-dipole forces.  
*Waterstofbindings is sterker as Londonkragte/Dispersiekragte/Geïnduseerde dipoolkragte en dipool-dipoolkragte.*
- More energy will be needed to overcome/break(inter-molecular) forces. ✓  
*Meer energie word benodig om (intermolekulêre) kragte te oorkom/te breek.* (3)

3.4.1 Organic molecules with the same molecular formula ✓ but different structural formulae. ✓  
*Organiese verbindings met dieselfde molekulêre formule maar verskillende struktuurformules.* (2)

3.4.2 CHAIN ✓/KETTING (1)

3.4.3 SMALLER THAN ✓/KLEINER AS



#### **STRUCTURE/STUKTUUR:**

Compound **D** are branched/more compact/more spherical/smaller contact area/smaller surface(over which intermolecular forces act.) ✓

*Verbinding **D** is meer vertak/meer kompak/meer series/kleiner kontak area/kleiner oppervlak(waaroor intermolekulêre kragte werk.)*


#### **INTERMOLECULAR FORCES/INTERMOLEKULÊRE KRAGTE**

Weaker/Less strength/Decrease in strength of Van der Waals forces/ London forces/Dispersion forces. ✓

*Swakker/Afname in sterkte van Van der Waalskragte/Londonkragte/ Dispersiekragte.*



4.4.1 (Concentrated) sulphuric acid ✓/hydrogen sulphate/H<sub>2</sub>SO<sub>4</sub>  
 Gekonsentreerde swaelsuur/waterstofsulfaat/ H<sub>2</sub>SO<sub>4</sub> (1)



**Marking criteria/Nasienglyne:**

- Whole structure correct/Hele struktuur korrek: 2/2
- Only functional group correct  
 Slegs funksionele groep korrek Max./Maks. 1/2


(2)

4.5.1

- Heat ✓/mild temperature over waterbath.  
Verhit/matige temperatuur oor 'n waterbad.
- Add concentrated sulphuric acid/H<sub>2</sub>SO<sub>4</sub> ✓  
Voeg gekonsentreerde swawelsuur/waterstofsulfaat/H<sub>2</sub>SO<sub>4</sub> by.

(2)

4.5.2



**Marking criteria/Nasienglyne:**

- Only functional group ✓  
 Slegs funksionele groep
- Whole structure correct ✓  
 Hele struktuur korrek 2/2

(2)

4.5.3 Butyl ✓ methanoate ✓/ butielmetanoaat (2)  
**[17]**

**QUESTION/VRAAG 5**

5.1.1 Use magnesium powder. ✓  
 Gebruik magnesiumpoeier. (1)

5.1.2 Increase concentration (H<sub>2</sub>SO<sub>4</sub>) ✓/Toename in konsentrasie (H<sub>2</sub>SO<sub>4</sub>) (1)

5.1.3 DECREASES ✓/VERLAAG (1)

5.1.4 NO EFFECT ✓/GEEN EFFEK (1)

5.2.1 Surface area ✓ (of Zn)/State of division(of Zn)  
*Oppervlaksarea(van Zn) /Toestand van verdeeldheid(van Zn)* (1)

5.2.2 Reaction stops ✓/come to completion/no more hydrogen gas is produced since zinc is used up. ✓  
*Reaksie stop/kom tot stilstand/geen waterstofgas word geproduseer want sink is opgebruik.* (2)

5.2.3 EQUAL TO ✓/GELYK AAN



The same Volume of H<sub>2</sub>(g) was produced. ✓  
*Dieselfde volume H<sub>2</sub>(g) is geproduseer.* (2)

5.2.4 (a) Average rate/Gemiddelde tempo =  $\Delta V/\Delta t$   
 $= (0,06 - 0) \checkmark / (30 - 0) \checkmark$   
 $= 2 \times 10^{-3} \checkmark (\text{mol} \cdot \text{dm}^{-3} \cdot \text{s}^{-1})$  or/of 0,002 (3)

(b)  $n = cV \checkmark = 0,4 \times 100/1000 \checkmark = 0,04 \text{ mol} \checkmark$  (3)

(c) **Marking criteria/Nasienriglyne:**

- Use 24,3 dm<sup>3</sup>.mol<sup>-1</sup> substituted in the correct formula. ✓  
*Gebruik 24,3 dm<sup>3</sup>.mol<sup>-1</sup> vervang in die korrekte formule.*
- Calculate n(HCl)<sub>reacted</sub> using the mol ratio 1 : 2. ✓  
*Bereken n(HCl)<sub>gereageer</sub> deur molverhouding 1 : 2 te gebruik.*
- Calculate n(HCl)<sub>reacted</sub> ✓  
*Bereken n(HCl)<sub>gereageer</sub>*
- Use 36,5 g.mol<sup>-1</sup> substitute in correct formule. ✓  
*Gebruik 36,5 g.mol<sup>-1</sup> vervang in die korrekte formule.*
- Finale antwoord/Final answer. (1,28–1,31 g) ✓



**POSITIVE MARKING from QUESTION 5.2.4 b**

**POSITIEWE NASIEN vanaf VRAAG 5.2.4 b**

$$n(\text{H}_2)_{\text{produced/berei}} = V/V_m = 0,06/24,3 \checkmark = 2,47 \times 10^{-3} \text{ mol} \quad (0,002)$$

$$n(\text{HCl})_{\text{reacted/gereageer}} = 2 \times 2,47 \times 10^{-3} \checkmark \quad \text{Ratio/Verhouding} \\ = 4,94 \times 10^{-3} \text{ mol} \quad (0,004)$$

$$n(\text{HCl})_{\text{remaining/oorgebly}} = n_{\text{initial/begin}} - n_{\text{produced/berei}} \\ = 0,04 - 4,94 \times 10^{-3} \checkmark \\ = 0,03506 \text{ mol} \quad (0,036)$$

$$m = nM = 0,03506 \times 36,5 \checkmark = 1,28 \text{ g} \checkmark \quad \text{Range/Gebied (1,28–1,314 g)} \quad (5)$$

5.2.5 The minimum energy required to start a chemical reaction. ✓✓  
*Die minimum energie benodig om 'n reaksie te begin.* (2 or/of 0) (2)

5.2.6 Experiment/Eksperiment III ✓ (1)

- 5.2.7
- A catalyst ✓ provides an alternative pathway of lower activation energy ( $E_A$ ). ✓  
*’n Katalisator verskaf ’n alternatiewe pad van laer aktiveringsenergie.*
  - More particles will have sufficient/enough kinetic energy ( $E_k$ ) to react. ✓  
*/More particles with  $E_k \geq E_A$ .  
Meer deeltjies het voldoende(genoegsame) kinetiese energie ( $E_k$ ) om te reageer./Meer deeltjies het  $E_k \geq E_A$ .*
  - More effective collisions per unit time/second. ✓  
*Meer effektiewe botsings per eenheidstyd/sekonde.*

**OR/OF**

Rate/frequency of effective collisions increases.

*Tempo/frekwensie van effektiewe botsings neem toe.*

(4)

[27]

### QUESTION/VRAAG 6

- 6.1 Stage at which the rate of forward reaction equals the rate of reverse. ✓✓  
*Die toestand/stadium in ’n chemiese reaksie wanneer die tempo van die voorwaartse reaksie gelyk is aan die tempo van die terugwaartse reaksie.*

**OR/OF**

The stage where the concentrations/quantities of reactants and products remain constant.

*Die toestand wanneer die konsentrasies/hoeveelhede van reaktanse en produkte konstant bly.*

**(2 or/of 0)**

(2)

- 6.2 Closed system ✓/Geslote sisteem  
Reversible reaction ✓/Omkeerbare reaksie.

(2)

6.3.1 NO EFFECT ✓/GEEN EFFEK

(1)

6.3.2 INCREASES ✓/TOENEEM

(1)

6.4.1 REVERSE ✓/TERUGWAARTSE

(1)

- 6.4.2 Increase in pressure ✓ /Decrease volume; Addition of a catalyst. ✓  
*Verhoog druk/Afname in volume; Byvoeging van ’n katalisator.*

(2)

- 6.5.1 LOW (yield) ✓/LAE (opbrengs)  
 $K_c$  is low. ✓/ $K_c$  is laag.

(2)

## 6.5.2

**Marking Criteria/Nasienriglyne:**

- Divide by 34 to calculate  $n(\text{H}_2\text{S})_{\text{equilibrium}}$ . ✓  
*Verdeel deur 34 om  $n(\text{H}_2\text{S})_{\text{ewewig}}$  te bereken.*
- Use mole ratio  $\text{H}_2:\text{H}_2\text{S}$ /Gebruik mol verhouding  $\text{H}_2:\text{H}_2\text{S}$  ✓ (1 : 1)
- Divide  $n(\text{H}_2\text{S})_{\text{equilibrium}}$  by V./Verdeel  $n(\text{H}_2\text{S})_{\text{ewewig}}$  deur V. ✓
- Correct  $K_c$  expression/Korrekte  $K_c$  uitdrukking. ✓
- Substitution of  $K_c$  -value./Vervanging van  $K_c$  waarde. ✓
- Substitution into  $K_c$  expression/Substutueer in  $K_c$  uitdrukking. ✓
- Calculate /Bereken  $n(\text{H}_2)_{\text{equilibrium/ewewig}}$  ✓
- Final answer/Finale antwoord  $n(\text{H}_2)_{\text{initial/begin}} = 2,45 \text{ mol}$  ✓

**OPTION/OPSIE 1**

$$n(\text{H}_2\text{S})_{\text{equilibrium/ewewig}} = m/M = 17/34 \checkmark = 0,5 \text{ mol}$$

	$\text{H}_2$	S	$\text{H}_2\text{S}$	
$n_{\text{initial/begin}}$ (mol)			0	ratio/ verhouding ✓
$\Delta n$ (mol)	0,5	0,5	0,5	
$n_{\text{equilibrium/ewewig}}$ (mol)			0,5	
$C_{\text{equilibrium/ewewig}}$ (mol.dm <sup>-3</sup> )	$n_{\text{equilibrium/ewewig}}/V$		$0,5/V$	$\div V$ ✓

$$K_c = [\text{H}_2\text{S}]/[\text{H}_2] \checkmark$$

$$2,56 \times 10^{-1} \checkmark = (0,5/V)/(n(\text{H}_2)_{\text{equilibrium/ewewig}}/V) \checkmark$$

$$n(\text{H}_2)_{\text{equilibrium/ewewig}} = 1,95 \text{ mol} \checkmark$$

$$n(\text{H}_2)_{\text{initial/begin}} = 0,5 + 1,95 = 2.45 \text{ mol} \checkmark$$

(8)

## 6.5.3

**POSITIVE MARKING from QUESTION 6.5.2****POSITIEWE NASIEN vanaf VRAAG 6.5.2**

$$90/100 n_i(\text{S}) \checkmark = \frac{n_i(\text{S}) - 0,5}{n_i(\text{S})} \checkmark$$

$$n_i(\text{S}) = 5 \text{ mol} \checkmark$$

(3)

## 6.5.4

**Graph/Grafiek Q ✓**

- As temperature increases,  $K_c$  decreases. ✓  
*Indien die temperatuur toeneem, neem  $K_c$  af.*
- $[\text{H}_2\text{S}]$  decreases ✓ /  $[\text{H}_2]$  increases.  
 *$[\text{H}_2\text{S}]$  neem af /  $[\text{H}_2]$  verhoog.*
- Reverse reaction is favoured by an increase in temperature. ✓  
*Terugwaartse reaksie word bevoordeel deur 'n verhoging in temperatuur.*

(4)

**[26]**

## QUESTION/VRAAG 7

- 7.1.1 Hydrolysis ✓/Hidrolise (1)
- 7.1.2 Transfer of proton ✓( $H^+$ ) occurs./  $CO_3^{2-}$  gains a proton /  $H_2O$  loses a proton.  
Oordrag van proton( $H^+$ ) vind plaas/ $CO_3^{2-}$  ontvang 'n proton/ $H_2O$  verloor 'n proton( $H^+$ ). (1)
- 7.1.3  $OH^-$  ✓ (1)
- 7.1.4  $K_b < 1 \times 10^{-14}$  ✓ (1)
- 7.1.5 Substance that can act either as an acid or base./ ✓✓  
'n Stof wat as beide suur of basis kan optree. (2)
- 7.1.6  $H_2CO_3$  ✓ (1)
- 7.1.7  $CO_3^{2-}$  ✓ (1)
- 7.2.1 An acid that donates TWO protons/ $H^+$ / $H_3O^+$ -ions. ✓✓  
'n Suur wat TWEE protone/ $H^+$ / $H_3O^+$ -ione vrystel. (2)
- 7.2.2  $pH = -\log [H_3O^+]$  ✓  
 $1,3$  ✓ =  $-\log [H_3O^+]$   
 $[H_3O^+] = 10^{-1,3}$   
 $[OH^-][H_3O^+] = 1 \times 10^{-14}$   
 $[OH^-] \times 10^{-1,3} = 1 \times 10^{-14}$  ✓  
 $[OH^-] = 10^{-12,7} \text{ mol.dm}^{-3}$  ✓ =  $1,995 \times 10^{-13} \text{ mol.dm}^{-3}$  (4)

### 7.3.1 POSITIVE MARKING FROM Q 7.2.2/POSITIEWE NASIEN VANAF V7.2.2 OPTION /OPSIE 1



$$[H_3O^+] = 10^{-1,3} \text{ mol.dm}^{-3}$$

$$[\text{Acid}] = \frac{1}{2} \times 10^{-1,3} \text{ ✓} = 0,0251 \text{ mol.dm}^{-3}$$

$$\text{Dilution } c_1V_1 = c_2V_2$$

$$0,0251(8) = c_2 \times 100 \text{ ✓}$$

$$c_2(\text{dilute}) = 2,008 \times 10^{-3} \text{ mol.dm}^{-3}$$

$$n(\text{acid reacting}) = cV = 2,008 \times 10^{-3} \times 25/1000 \text{ ✓} = 5,02 \times 10^{-5} \text{ mol}$$

$$n(\text{base reacting}) = 2 \times 5,02 \times 10^{-5} \text{ ✓} = 1,004 \times 10^{-4} \text{ mol}$$

$$c(\text{base}) = n/V = 1,004 \times 10^{-4} / 14,2 \times 10^{-3} \text{ ✓} = 7,07 \times 10^{-3} \text{ mol.dm}^{-3} \text{ ✓} \quad (6)$$

## OPTION/OPSIE 2

$$[\text{H}_3\text{O}^+] = 10^{-1,3} \text{ mol.dm}^{-3}$$

$$[\text{Acid}] = \frac{1}{2} \times 10^{-1,3} \checkmark = 0,0251 \text{ mol.dm}^{-3}$$

$$c_1V_1 = c_2V_2$$

$$(0,0251)(8) = c_2 \times 100 \checkmark$$

$$c_2 = 2,008 \times 10^{-3} \text{ mol.dm}^{-3}$$

$$\frac{c_a V_a}{c_b V_b} = \frac{n_a}{n_b}$$

$$\frac{c_b V_b}{c_a V_a} = \frac{n_b}{n_a}$$

$$\frac{2,008 \times 10^{-3} (25) \checkmark}{c_b \times 14,2 \checkmark} = \frac{1 \checkmark}{2}$$

$$c_b = 7,07 \times 10^{-3} \text{ mol.dm}^{-3} \checkmark$$

**Range/Gebied** ( $7,04 \times 10^{-3}$  to/tot  $7,07 \times 10^{-3} \text{ mol.dm}^{-3}$ )

7.3.2 **B**✓



Titration of a strong base and a strong acid ✓ (solution at end point neutral.)

*Titrasie van 'n sterk basis en 'n sterk suur (oplossing is neutraal by eindpunt.)*

(2)

[22]

**TOTAL/TOTAAL: 150**